

# CALCULUS

(1)

1. Limits and Continuity

2. Differentiation

3. Definite Integrals

    Improper Integrals

4. Partial Differentiation

5. Multiple Integrals

6. Vector Differentiation

7. Vector Integration

8. Fourier Series

Calculus is defined as science of acceleration, retardation and variation.

function: $\rightarrow$  A relation between two sets 'A' & 'B' if  $x \in A$   $\exists$  a unique  $y \in B$  s.t.  $f(x) = y$

(i) Explicit function: $\rightarrow z = f(x_1, x_2, \dots, x_n)$   
                                ↑                                 ↑  
                                Dependent              Independent  
                                variable                 variable

(ii) Implicit function: $\rightarrow \phi(z, x_1, x_2, \dots, x_n) = C$

(iii) Composite function: $\rightarrow$  If  $z = f(x, y)$ , where  $x = \phi(t)$  &  $y = \psi(t)$ ,  
i.e.  $z$  is function of some function.


### Some Special functions

(i) Even function: $\rightarrow f(-x) = f(x)$  Eg:-  $\cos x$ ,  $|x|$ , ...

(ii) Odd function: $\rightarrow f(-x) = -f(x)$  Eg:-  $\sin x$ ,  $x$ , ...

(iii) Modulus function: $\rightarrow$

$$f(x) = |x| = \begin{cases} x & ; x > 0 \\ -x & ; x < 0 \\ 0 & ; x = 0 \end{cases}$$




$$\frac{d}{dx} |x| = \frac{|x|}{x} \text{ for } x \neq 0$$

(iv) Step-function/Greatest Integer function

$$f(x) = [x] = n \in \mathbb{Z}$$

where,  $n \leq x < n+1$

$$\text{Eg:- } [7.2] = 7 ; [7.999] = 7 ; [-1.2] = -2$$



### Symmetric Properties of the curve: $\rightarrow$

Let  $f(x, y) = C$  be the eqn of the curve

(i) If  $f(x, y)$  contains only even powers of  $x$  i.e.  $\text{even}$

$f(-x, y) = f(x, y)$  then it is symmetric about  $y$ -axis.

(ii) If  $f(x, y)$  contains only even powers of  $y$  i.e.  $f(x, -y) = f(x, y)$

then it is symmetric about  $x$ -axis.

(iii) If  $f(x, y) = f(y, x)$ , then, the curve is symmetric about  $y=x$ .

2

1. Limit of a function: $\rightarrow$  Let  $f(x)$  be defined in deleted neighbourhood of  $a \in \mathbb{R}$ , then,  $l \in \mathbb{R}$  is said to be limit of  $f(x)$  as  $x$  approaches  $a$  if for given  $\epsilon > 0 \exists \delta > 0$  such that  $|f(x) - l| < \epsilon$  whenever  $|x-a| < \delta$ .

$$\boxed{\lim_{x \rightarrow a} f(x) = l}$$

Left limit: $\rightarrow$  when  $x < a, x \rightarrow a$

$$\lim_{x \rightarrow a^-} f(x) = \lim_{h \rightarrow 0} f(a-h)$$

Right Limit: $\rightarrow$  when  $x > a, x \rightarrow a$

$$\lim_{x \rightarrow a^+} f(x) = \lim_{h \rightarrow 0} f(a+h)$$

A limit exists iff  $LHL = RHL$

Indeterminate form: $\rightarrow \frac{0}{0}, \frac{\infty}{\infty}, 0 \times \infty, \infty - \infty, 0^0, 1^\infty, \infty^0$

Whenever we have  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$  [as  $\frac{0}{0}$  or  $\frac{\infty}{\infty}$ ] =  $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$

This rule is applied until we are free from indeterminate form. (This rule is called L'Hospital Rule)

### Standard Limits:-

$$(i) \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n a^{n-1}$$

$$(ii) \lim_{x \rightarrow 0} \frac{e^{mx} - 1}{x} = m$$

$$(iii) \lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log a$$

$$(iv) \lim_{x \rightarrow 0} [1 + ax]^{1/x} = e^a$$

$$(v) \lim_{x \rightarrow \infty} \left[1 + \frac{a}{x}\right]^x = e^a$$

$$(vi) \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad \text{or} \quad \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$$

$$(vii) \lim_{x \rightarrow 0} \frac{\sin x}{x} = 0$$

$$(viii) \lim_{x \rightarrow 0} \frac{\sin mx}{x} = m$$

$$(ix) \lim_{x \rightarrow 0} \left[ \frac{a^x + b^x}{2} \right]^{1/x} = \sqrt{ab}$$

$$(x) \lim_{x \rightarrow 0} [\cos x + a \sin bx]^{1/x} = e^{ab}$$

$$(xi) \lim_{x \rightarrow 0} \left[ \frac{1 - \cos ax}{x^2} \right] = \frac{a^2}{2}$$

Questions: →

3

$$1. \lim_{x \rightarrow 0} \frac{1 - \cos 3x}{x \sin 2x}$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow 0} \frac{1 - \cos 3x}{x \sin 2x} = \lim_{x \rightarrow 0} \frac{3 \sin 3x}{\sin 2x + 2x \cos 2x}$$

$$= \lim_{x \rightarrow 0} \frac{9 \cos 3x}{2 \cos 2x + 2 \cos 2x - 4x \sin 2x}$$

$$= \frac{9}{4}$$

OR

$$\lim_{x \rightarrow 0} \frac{(1 - \cos 3x)/x^2}{(\sin 2x)/x} = \frac{3^2/2}{2} = \frac{9}{4}$$

$$2. \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin^2 2x}{(x - \frac{\pi}{2})^2} \text{ is } \dots$$

- (a) 1      (b) 2      (c) 4      (d) -4

$$\text{Soln:} \rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{\left[ \sin 2\left(\frac{\pi}{2} - x\right) \right]^2}{\left(\frac{\pi}{2} - x\right)^2}$$

Take,  $\frac{\pi}{2} - x = t$ , then

$$\lim_{t \rightarrow 0} \left[ \frac{\sin 2t}{t} \right]^2 = (2)^2 = 4$$

$$3. \lim_{x \rightarrow 0} \frac{\log x}{\cot x}$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow 0} \frac{\log x}{\cot x} = \lim_{x \rightarrow 0} \frac{\frac{1}{x}}{-\operatorname{cosec}^2 x} = - \left[ \lim_{x \rightarrow 0} \left( \frac{\sin x}{x} \cdot \sin x \right) \right]$$

$$= -1 \times 0 = 0$$

$$4. \lim_{x \rightarrow \frac{\pi}{2}} \frac{\tan x}{\log(\cos x)}$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{\tan x}{\log(\cos x)} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sec^2 x}{-\sec x \sin x} = - \lim_{x \rightarrow \frac{\pi}{2}} \frac{1}{\sin x \cos x} \\ = -\infty.$$

$$5. \lim_{x \rightarrow \frac{\pi}{2}} \tan x$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \tan x = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\sin x}{\cos x} \quad \text{Limit doesn't exist} \quad (\because \text{LHL} \neq \text{RHL})$$

$$6. \lim_{x \rightarrow 0} \sin x \log x^2$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow 0} \sin x \log x^2 = \lim_{x \rightarrow 0} \frac{\log x^2}{\csc x} = \lim_{x \rightarrow 0} \frac{2/x}{-\cosec x \cot x} \\ = \lim_{x \rightarrow 0} \left( -2 \times \frac{\sin x}{x} \times \tan x \right) \\ = -2 \times 1 \times 0 = 0.$$

$$\text{Note:} \rightarrow \log a^0 = \begin{cases} \infty & ; a < 1 \\ -\infty & ; a > 1 \end{cases}$$

$$7. \lim_{x \rightarrow 1} (x-1) \tan \frac{\pi x}{2}$$

$$\text{Soln:} \rightarrow \lim_{x \rightarrow 1} (x-1) \tan \frac{\pi x}{2} = \lim_{x \rightarrow 1} \frac{(x-1)}{\cot \frac{\pi x}{2}} = \lim_{x \rightarrow 1} \left[ \frac{1}{-\frac{\pi}{2} \cosec^2 \left( \frac{\pi x}{2} \right)} \right] \\ = -\frac{2}{\pi}.$$

$$\text{Note:} \rightarrow (i) \lim_{x \rightarrow 0} x \sin(1/x) = 0$$

$$(iii) \lim_{x \rightarrow 0} \sin(1/x) \text{ does not}$$

exists.

$$(ii) \lim_{x \rightarrow \infty} x \sin(1/x) = 1$$

8.  $\lim_{x \rightarrow 0} \left[ \frac{1}{x} - \frac{1}{x^2} \log(1+x) \right]$  is

4

- (a)  $\frac{1}{2}$     (b)  $\frac{1}{3}$     (c)  $-\frac{1}{2}$     (d)  $-\frac{1}{3}$

$$\text{Soln: } \lim_{x \rightarrow 0} \left[ \frac{x - \log(1+x)}{x^2} \right] = \lim_{x \rightarrow 0} \left[ \frac{1 - \frac{1}{1+x}}{\frac{2x}{2x}} \right]$$

$$= \lim_{x \rightarrow 0} \frac{\frac{1}{(1+x)^2}}{2} = \frac{1}{2}$$

9.  $\lim_{x \rightarrow 0} \left[ \frac{1}{x^2} - \frac{1}{\sin^2 x} \right]$  is

- (a)  $\frac{1}{3}$     (b)  $-\frac{1}{3}$     (c)  $\frac{1}{2}$     (d)  $\frac{1}{4}$

$$\text{Soln: } \lim_{x \rightarrow 0} \left[ \frac{\sin^2 x - x^2}{x^2 \sin^2 x} \right] = \lim_{x \rightarrow 0} \left[ \frac{\sin^2 x - x^2}{x^4 \cdot \cancel{\sin^2 x}} \right]$$

$$= \lim_{x \rightarrow 0} \left[ \frac{\sin^2 x - x^2}{x^4} \right]$$

$$= \lim_{x \rightarrow 0} \left[ \frac{\sin 2x - 2x}{4x^3} \right]$$

$$= \lim_{x \rightarrow 0} \frac{2 \cos 2x - 2}{12x^2}$$

$$= -\frac{2}{12} \lim_{x \rightarrow 0} \left( \frac{1 - \cos 2x}{x^2} \right)$$

$$= -\frac{2}{12} \times \frac{4}{2} = -\frac{1}{3}$$

10.  $\lim_{x \rightarrow 0} x^x$ .

Soln: Let  $y = x^x$

$$\Rightarrow \log y = x \log x$$

$$\Rightarrow \lim_{x \rightarrow 0} [\log y] = \lim_{x \rightarrow 0} (x \log x) = \log (\lim_{x \rightarrow 0} x^x) = 0$$

$$11. \lim_{x \rightarrow \frac{\pi}{2}} (\cos x)^{\frac{1}{\sin x}}$$

$$12. \lim_{x \rightarrow 1} [\log x]^{\log x}$$

Note: → If we have  $1^\infty$  form,  $\lim_{x \rightarrow a} [f(x)]^{g(x)} = e^{\lim_{x \rightarrow a} g(x)[f(x)-1]}$

$$13. \lim_{x \rightarrow 0} (1 - \sin x)^{\frac{1}{\sin x}}$$

$$\lim_{x \rightarrow 0} \frac{1}{\sin x} [1 - \sin x - 1]$$

$$\text{Soln:} \rightarrow e^{\lim_{x \rightarrow 0} \frac{1}{\sin x} [1 - \sin x - 1]} = e^{-1} = \frac{1}{e}$$

$$14. \lim_{x \rightarrow 0} (\cos x)^{\frac{1}{x^2}}$$

$$= e^{-\lim_{x \rightarrow 0} \left[ \frac{1 - \cos x}{x^2} \right]} = e^{-\frac{1}{2}}$$

$$\text{Soln:} \rightarrow e^{\lim_{x \rightarrow 0} \frac{1}{x^2} [\cos x - 1]} = e^{-\frac{1}{2}} = \frac{1}{\sqrt{e}}$$

$$15. \lim_{x \rightarrow 0} \left( \frac{2^x + 8^x}{2} \right)^{\frac{1}{x}} = \sqrt{ab} = \sqrt{2 \times 8} = \sqrt{16} = 4.$$

$$16. \lim_{x \rightarrow 0} \left( \frac{2^x + 4^x + 8^x}{3} \right)^{\frac{1}{x}} = \sqrt[3]{abc} = \sqrt[3]{2 \times 4 \times 8} = 4.$$

$$17. \lim_{x \rightarrow 0} [\cos x + 2 \sin 3x]^{\frac{1}{x}} = e^{ab} = e^{2 \times 3} = e^6.$$

18.  $\lim_{x \rightarrow 0} [2 \cos x + 3 \sin 4x]^{1/x}$

5

$$= \lim_{x \rightarrow 0} 2^{\frac{1}{x}} \left[ \cos x + \frac{3}{2} \sin 4x \right]^{\frac{1}{x}} = \lim_{x \rightarrow 0} 2^{\frac{1}{x}} \cdot \lim_{x \rightarrow 0} \left[ \cos x + \frac{3}{2} \sin 4x \right]^{\frac{1}{x}}$$

= does not exist. ( $\because \text{LHL} \neq \text{RHL}$  for  $\lim_{x \rightarrow 0} 2^{\frac{1}{x}}$ )

19.  $\lim_{x \rightarrow 0} \frac{|x|}{x} = ?$

Sol<sup>n</sup>:— LHL =  $\lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{h \rightarrow 0} \frac{|h|}{-h} = -1$

RHL =  $\lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{h \rightarrow 0} \frac{|h|}{h} = 1$

$\therefore \text{LHL} \neq \text{RHL}$

$\therefore$  limit doesn't exist

20.  $\lim_{x \rightarrow a} [x]$  doesn't exist when  $a$  is \_\_\_\_\_

- (a) Real no.
- (b) Rational no.
- (c) Integer
- (d) all of these

Sol<sup>n</sup>:— Let  $a = 2$

LHL =  $\lim_{x \rightarrow 2^-} [x] = 1$

RHL =  $\lim_{x \rightarrow 2^+} [x] = 2$

$\therefore \text{LHL} \neq \text{RHL} \Rightarrow$  limit doesn't exist.

### Continuity of a function:

(i) Continuity at a point: → A fun is said to be continuous at a point  $x=a$ , if  $\lim_{x \rightarrow a} f(x) = f(a)$

(ii) Continuity in an interval: → A fun  $f(x)$  is said to be continuous in  $[a, b]$  if it satisfies the following three conditions:—

(a)  $f(x)$  is continuous  $\forall x \in (a, b)$

(b)  $\lim_{x \rightarrow a^+} f(x) = f(a)$

(c)  $\lim_{x \rightarrow b^-} f(x) = f(b)$

$$\text{Ex:- (i) If } f(x) = \begin{cases} \frac{x^2-4}{x-2} & \text{if } x \neq 2 \\ 2 & \text{if } x=2 \end{cases} \quad \text{check its continuity}$$

at  $x=2$ .

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \left( \frac{x^2-4}{x-2} \right) = \lim_{x \rightarrow 2} \left( \frac{(x-2)(x+2)}{x-2} \right) = \lim_{x \rightarrow 2} (x+2) = 4$$

But at  $x=2$ ,  $f(x)=2$

∴  $f(x)$  is not continuous at  $x=2$ .

$$(ii) \text{ If } f(x) = \begin{cases} (1+3x)^{\frac{1}{x}} & ; x \neq 0 \\ e^x & ; x=0 \end{cases}$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (1+3x)^{\frac{1}{x}} = e^{\lim_{x \rightarrow 0} \frac{1}{x} (1+3x-1)} = e^{3} = e$$

∴  $f(x)$  is continuous at  $x=0$ .

Questions:-

Ques. 1. If  $f(x) = \begin{cases} 0 & ; x=0 \\ \frac{1}{2}-x & ; 0 < x < \frac{1}{2} \\ \frac{1}{2} & ; x=\frac{1}{2} \\ \frac{3}{2}-x & ; \frac{1}{2} < x < 1 \\ 1 & ; x \geq 1 \end{cases}$

6

Then which of the following is true :-

- (a)  $f(x)$  is right continuous at  $x=0$
- (b)  $f(x)$  is discontinuous at  $x=\frac{1}{2}$
- (c)  $f(x)$  is continuous at  $x=1$
- (d) b & c

Soln:- (a)  $f(0)=0$

2. Differentiation: $\rightarrow$  A fun  $f(x)$  is said to be differentiable at a pt.

$x=c$ , if  $\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$  exists & finite. & is represented by  $f'(c)$ .

Left Hand Derivative: $\rightarrow \lim_{h \rightarrow 0} \frac{f(c-h) - f(c)}{-h}$

Right Hand Derivative: $\rightarrow \lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}$

Necessary cond<sup>n</sup> for a fun to be differentiable is  $LHD = RHD$ .

Note: $\rightarrow$  (i)  $f(x) = |x|$  is not differentiable at  $x=0$

$$LHD = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{|-h| - 0}{-h} = -1$$

$$RHD = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{|h| - 0}{h} = 1$$

$$\therefore LHD \neq RHD$$

$\Rightarrow$  Not differentiable

(ii)  $|x-a|$  is not differentiable at  $x=a$ .

(iii)  $|ax+b|$  is not differentiable at  $x=-b/a$ .

Questions: $\rightarrow$

1.  $f(x) = |x| + |x+1| + |x-2|$  is differentiable at  $x = \underline{\hspace{2cm}}$ .

- (a) 0      (b) 1      (c) -1      (d) 2

2. Let  $f(x) = |x+1|$  be defined in the interval  $[0, 4]$  then,

(a)  $f(x)$  is continuous & differentiable

(b)  $f(x)$  is continuous but non-differentiable

(c)  $f(x)$  is not continuous but differentiable

(d)  $f(x)$  is neither differentiable nor continuous

3. If  $f(x) = |x|^3$  where  $x \in \mathbb{R}$ , then,  $f(x)$  at  $x=0$  is \_\_\_\_\_ 7

- (a) continuous but not differential  
(b) Once differentiable but not twice  
 (c) Twice differentiable but not thrice  
(d) Thrice differentiable

Soln:-

$$f(x) = |x|^3 = \begin{cases} x^3 & ; x > 0 \\ -x^3 & ; x < 0 \\ 0 & ; x = 0 \end{cases}$$

$f(0) = 0$ , LHL = 0, RHL = 0  $\Rightarrow$  continuous

$$\text{LHD} = \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0^-} \frac{|h|^3 - 0}{-h} = 0$$

$$\text{RHD} = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{|h|^3 - 0}{h} = 0$$

$\therefore \text{LHD} = \text{RHD} \Rightarrow$  differentiable

$$f'(x) = \begin{cases} 3x^2 & ; x > 0 \\ -3x^2 & ; x < 0 \\ 0 & ; x = 0 \end{cases}$$

$$f''(x) = \begin{cases} 6x & ; x > 0 \\ -6x & ; x < 0 \\ 0 & ; x = 0 \end{cases} = 6|x|$$

$$f'''(x) = \begin{cases} 6 & ; x > 0 \\ -6 & ; x < 0 \\ \text{not diff.} & ; x = 0 \end{cases}$$

4. If  $f(x) = \begin{cases} 2+x & x \geq 0 \\ 2-x & x < 0 \end{cases}$  then  $f(x)$  at  $x=0$  is \_\_\_\_\_

- (a) Continuous & differentiable
- (b) continuous but not differentiable
- (c) Differentiable but not continuous
- (d) Neither diff. nor continuous

Soln.  $\rightarrow f(0) = 2+0 = 2, LHL = 2-0 = 2, RHL = 2+0 = 2 \Rightarrow$  continuous

$$LHD = \lim_{h \rightarrow 0} \frac{[2-(-h)]-2}{-h} = -1 \quad \left. \right\} \text{Non-Differentiable.}$$

$$RHD = \lim_{h \rightarrow 0} \frac{[2+h-2]}{h} = 1 \quad \left. \right\}$$

Note:  $\rightarrow$  (i) Every differentiable fun is a continuous fun.

(ii) But every continuous fun is not differentiable.

### Mean-Value Theorem:

(i) Rolle's Theorem:  $\rightarrow$  Let  $f(x)$  be defined in  $[a, b]$  s.t. it satisfies

three condn:-

(a)  $f(x)$  is continuous fun in  $[a, b]$

(b)  $f(x)$  is differentiable fun in  $(a, b)$

(c)  $f(a) = f(b)$

then, there exists atleast one point  $c \in (a, b)$  s.t.

$$f'(c) = 0$$

(ii) Lagrange's Mean Value Theorem:  $\rightarrow$  Let  $f(x)$  be defined in  $[a, b]$

s.t. it satisfies two condn:-

(a)  $f(x)$  is continuous fun in  $[a, b]$

(b)  $f(x)$  is differentiable fun in  $(a, b)$

then,  $\exists$  at least one point  $c$ , in  $(a, b)$  s.t.  $f'(c) = \frac{f(b) - f(a)}{b - a}$

Note:  $\rightarrow$  If  $f(x)$  is defined in  $[a, a+h]$  s.t.

(a)  $f(x)$  is continuous in  $[a, a+h]$

(b)  $f(x)$  is differentiable in  $(a, a+h)$

then  $\exists \theta \in (0, 1)$  s.t.

$$f(a+h) = f(a) + h f'(a+\theta h)$$

$$\theta = \frac{c-a}{a-b}$$

Questions:  $\rightarrow$

1. The mean-value 'c' for the fun.  $f(x) = e^x (\sin x - \cos x)$  in  $[\frac{\pi}{4}, \frac{5\pi}{4}]$

is \_\_\_\_\_

(a) 0

(b)  $\frac{\pi}{3}$

(c)  $\frac{\pi}{2}$

(d)  $\pi$

Sol<sup>n</sup>:  $\rightarrow f(\frac{\pi}{4}) = e^{\frac{\pi}{4}} (\sin \frac{\pi}{4} - \cos \frac{\pi}{4}) = 0$

$$f(\frac{5\pi}{4}) = e^{\frac{5\pi}{4}} (\sin \frac{5\pi}{4} - \cos \frac{5\pi}{4}) = 0$$

$$f'(x) = e^x [\cos x + \sin x] + e^x [\sin x - \cos x] = 2e^x \sin x$$

$$\Rightarrow f'(c) = 2e^c \sin c = 0$$

$$\Rightarrow c = 0, \pm \pi, \pm 2\pi, \dots$$

2. The mean-value 'c' for the fun.  $f(x) = x^3 - 6x^2 + 11x - 6$  in  $[0, 4]$  is

(a)  $2 + \frac{2}{\sqrt{3}}$

(b)  $2 - \frac{2}{\sqrt{3}}$

(c)  $2 \pm \frac{2}{\sqrt{3}}$

(d) None

Sol<sup>n</sup>:  $\rightarrow f(0) = -6$

$$f(4) = 64 - 96 + 44 - 6 = 6$$

$$f'(x) = 3x^2 - 12x + 11$$

$$\therefore f'(c) = 3c^2 - 12c + 11 = \frac{f(4) - f(0)}{4 - 0}$$

$$\Rightarrow 12c^2 - 48c + 44 = 12$$

$$\Rightarrow c = 2 \pm \frac{2}{\sqrt{3}}$$

3. The value of  $\zeta$  of  $f(b) - f(a) = (b-a)f'(\zeta)$  for the fun<sup>n</sup>  
 $f(x) = Ax^2 + Bx + C$ , in  $[a, b]$  is \_\_\_\_\_.

(a)  $\frac{b+a}{2}$       (b)  $\frac{b-a}{2}$       (c)  $b-a$       (d)  $b+a$

Sol<sup>n</sup>:  $\rightarrow f'(x) = 2Ax + B$

$$f'(\zeta) = \frac{f(b) - f(a)}{b-a}$$

$$\Rightarrow 2A\zeta + B = \frac{f(b) - f(a)}{b-a} = \frac{(Ab^2 + Bb + C) - (Aa^2 + Ba + C)}{b-a}$$

$$\Rightarrow 2A\zeta + B = (b+a)A + B$$

$$\Rightarrow \zeta = \frac{b+a}{2}$$

4. The mean-value 'c' for the fun<sup>n</sup>  $f(x) = 3x^2 + 5x + 11$  in the  $\left[\frac{11}{2}, \frac{16}{2}\right]$  is \_\_\_\_\_.

Sol<sup>n</sup>:  $c = \frac{\frac{11}{2} + \frac{16}{2}}{2} = \frac{27}{4}$

Note:  $\rightarrow$  If the fun<sup>n</sup>  $f(x)$  is polynomial of degree 2 i.e. quadratic

then,  $c$  will be the average value of the extreme value of  
 the fun<sup>n</sup> in given interval i.e.  $c = \frac{a+b}{2}$  if  $f(x)$  is defined in  $[a, b]$ .

5. The value of  $\theta \in (0, 1)$  for the fun<sup>n</sup>  $f(x) = \log x$  in  $[1, e]$  using  
 an appropriate mean-value theorem is \_\_\_\_\_.

Sol<sup>n</sup>:  $\rightarrow f'(x) = \frac{1}{x} \Rightarrow f'(c) = \frac{1}{c}$

$$f(1) = \log 1 = 0$$

$$f(e) = \log e = 1$$

$$\therefore \text{Using LMVT, } \frac{1}{c} = \frac{1-0}{e-1} \Rightarrow c = e-1$$

$$\therefore \theta = \frac{c-a}{b-a} = \frac{e-1-1}{e-1} = \frac{e-2}{e-1} \in (0, 1)$$

6. LMVT cannot be applied for  $f(x) = x^{1/3}$  in  $[-1, 1]$  because

- (a)  $f(x)$  is not continuous in  $[-1, 1]$
- (b)  $f(x)$  is not differentiable in  $(-1, 1)$
- (c) a & b
- (d)  $f(-1) \neq f(1)$

Sol<sup>n</sup>:  $f'(x) = \frac{1}{3}x^{-2/3} = \frac{1}{3x^{2/3}}$

$\therefore f(x)$  is not differentiable at  $x=0 \in (-1, 1)$

$$f(0) = 0$$

$$\text{LHL} = \lim_{h \rightarrow 0} f(0-h) = \lim_{h \rightarrow 0} (-h)^{1/3} = 0$$

$$\text{RHL} = \lim_{h \rightarrow 0} f(0+h) = \lim_{h \rightarrow 0} h^{1/3} = 0.$$

$$\text{LHL} = \text{RHL} = f(0) = 0 \Rightarrow \text{continuous.}$$

7. Rolle's Theorem cannot be applied for  $f(x) = |x+2|$  in  $[-2, 0]$ .

- (a)  $f(x)$  is not continuous in  $[-2, 0]$

- (b)  $f(x)$  is not differentiable in  $(-2, 0)$

- (c)  $f(-2) \neq f(0)$

- (d) b & c

8. If  $f'(x) = \frac{1}{5-x^2}$  and  $f(0) = 1$  then the lower & upper bounds of  $f(1)$  are \_\_\_\_\_.

Sol<sup>n</sup>: Let  $f(x)$  be defined in  $[0, 1]$ .

By LMVT,  $\exists c \in (0, 1)$  st.

$$f'(c) = \frac{f(1) - f(0)}{1-0}$$

$$\Rightarrow f'(c) = f(1) - 1$$

$$\min\{f'(x)\} < f'(c) < \max\{f'(x)\} \quad 0 \leq x \leq 1$$

$$\min\{f'(x)\} < f'(c) < \max\{f'(x)\}$$

$$\frac{1}{5} < f(1) - 1 < \frac{1}{4}$$

$$\Rightarrow 1 + \frac{1}{5} < f(1) < \frac{1}{4} + 1$$

(iii) Cauchy's Mean-Value Theorem: → Let  $f(x)$  &  $g(x)$  be defined in a closed interval  $[a, b]$  s.t. they satisfy the cond'':-

(a)  $f(x)$  &  $g(x)$  are continuous in  $[a, b]$

(b)  $f(x)$  &  $g(x)$  are differentiable in  $(a, b)$

(c)  $g'(x) \neq 0 \quad \forall x \in (a, b)$ , then,

$$\exists c \in (a, b) \text{ s.t. } \frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}$$

Questions:→

1. The mean-value  $c$  for the fun  $f(x) = e^x$  &  $g(x) = \bar{e}^x$  in  $[0, 1]$ .

is \_\_\_\_\_

$$\underline{\text{Sol'n}}: \rightarrow f'(x) = e^x \Rightarrow f'(c) = e^c$$

$$g'(x) = -\bar{e}^x \Rightarrow g'(c) = -\bar{e}^c$$

$$\therefore f(1) = e, \quad f(0) = 1$$

$$g(1) = \frac{1}{e}, \quad g(0) = 1$$

$$\therefore \frac{e^c}{-\bar{e}^{-c}} = \frac{e-1}{(1/e)-1} \Rightarrow -e^{2c} = \frac{[e-1]}{1-e} e$$

$$\Rightarrow +e^{2c} = e \Rightarrow c = \frac{1}{2} \in [0, 1].$$

2.  $f(x) = \sin x$  &  $g(x) = \cos x$  in  $[-\frac{\pi}{2}, 0]$  is \_\_\_\_\_.

Sol'n:→  $f'(x) = \cos x, \quad g'(x) = -\sin x \neq 0 \quad \forall x \in (-\frac{\pi}{2}, 0)$

$$\frac{f'(c)}{g'(c)} = \frac{f(0) - f(-\pi/2)}{g(0) - g(-\pi/2)}$$

$$\Rightarrow \frac{\cos c}{-\sin c} = \frac{0 - (-1)}{1 - 0} \Rightarrow -\cot c = 1 \Rightarrow \cot c = -1$$

$$\Rightarrow c = -\frac{\pi}{4} \in (-\frac{\pi}{2}, 0)$$
10

(iv) Taylor's Theorem:  $\rightarrow$  OR (Generalised Mean-Value Theorem)

Let  $f(x)$  be defined in  $[a, a+h]$ . s.t.

(a)  $f, f', f'', f''', \dots, f^{n-1}$  are continuous in  $[a, a+h]$

(b)  $f, f', f'', \dots, f^{n-1}$  are differentiable in  $(a, a+h)$

then  $\exists \theta \in (0, 1)$ , s.t.

$$f(a+h) = f(a) + h f'(a) + \frac{h^2}{2!} f''(a) + \dots + \frac{h^{n-1}}{(n-1)!} f^{n-1}(a) + R_n$$

$$\text{where } R_n = \frac{h^n}{(n-1)!} (1-\theta)^{n-p} f^n(a+\theta h)$$

for  $p=n \rightarrow$  Long Lagrange's form of remainder

$p=1 \rightarrow$  Cauchy's form of remainder

Case I: when  $p=n$ ,  $R_n = \frac{h^n}{n!} f^n(a+\theta h)$ .

Case II: when  $p=1$ ,  $R_n = \frac{h^n}{(n-1)!} (1-\theta)^{n-1} f^n(a+\theta h)$ .

Taylor's Series:  $\rightarrow$  As  $n \rightarrow \infty$ ,  $R_n \rightarrow 0$ , then

$$f(a+h) = f(a) + h f'(a) + \frac{h^2}{2!} f''(a) + \frac{h^3}{3!} f'''(a) + \dots$$

(i)  $f(x) = f(a) + (x-a) f'(a) + \frac{(x-a)^2}{2!} f''(a) + \dots$  is a Taylor's series expansion of  $f(x)$  about  $x=a$ .

(ii)  $f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \dots$  is a Taylor's series expansion of  $f(x)$  about  $x=0$ .

(Maclaurian's Series)

Questions: →

1. The coeff. of  $x^2$  in the Taylor's series expansion of  $\cos^2 x$  about  $x=0$  is —

- (a) 0      (b) 1      (c) -1      (d) 2

$$\text{Sol}^n: \rightarrow \text{coeff. of } x^2 = \frac{f''(0)}{2!} = \frac{-2}{2} = -1$$

$$\text{where, } f'(x) = -\sin 2x$$

$$f''(x) = -2\cos 2x \Rightarrow f''(0) = -2$$

2. The coeff. of  $(x-2)^4$  in the Taylor's series expansion of  $e^x$  about

$x=2$  is —

- (a)  $\frac{e^2}{2!}$       (b)  $\frac{e^2}{4!}$       (c)  $\frac{e^4}{2!}$       (d)  $\frac{e^4}{4!}$

$$\text{Sol}^n: \rightarrow \text{coeff. of } (x-2)^4 = \frac{f''''(2)}{4!} = \frac{e^2}{4!}$$

3. The coeff. of  $(x-\pi)^3$  in the power series expansion of  $e^x + \sin x$

in the ascending power of  $(x-\pi)$  is —

- (a)  $\frac{e^\pi}{6}$       (b)  $\frac{e^\pi+1}{3}$       (c)  $\frac{e^\pi-1}{3}$       (d) None

$$\text{Sol}^n: \rightarrow \text{coeff. of } (x-\pi)^3 = \frac{f^3(\pi)}{3!} = \frac{e^\pi+1}{6}$$

$$\text{where, } f'(x) = e^x + \cos x$$

$$f''(x) = e^x - \sin x$$

$$f'''(x) = e^x - \cos x \Rightarrow f^3(\pi) = e^\pi + 1$$

4. Which of the following fun ~~also~~ would have only odd powers of  $x$  in its Taylor's series expansion about  $x=0$ ,

- (a)  $\sin x^2$       (b)  $\cos x^2$       (c)  $\cos x^3$       (d)  $\sin x^3$

$$\text{Soln: } \cos t = 1 - \frac{t^2}{2!} + \frac{t^4}{4!} - \dots$$

$$\sin t = t - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots$$

11

5. The Taylor's Series expansion of  $f(x) = \tan^{-1}x$  about  $x=0$  is \_\_\_\_\_.

$$\text{Soln: } f(x) = f(0) + x f'(0) + \frac{x^2}{2!} f''(0) + \dots$$

$$f(x) = \tan^{-1}x$$

$$f(0) = \tan^{-1}0 = 0$$

$$f'(x) = \frac{1}{1+x^2} \Rightarrow f'(0) = 1$$

$$f''(x) = \frac{-2x}{(1+x^2)^2} \Rightarrow f''(0) = 0$$

$$f'''(x) = -2 \left[ \frac{(1+x^2)^2 + x \cdot 2(1+x^2) \cdot 2x}{(1+x^2)^4} \right] = -2 \left[ \frac{1+x^2 - 4x^3}{(1+x^2)^3} \right] = -2 \left[ \frac{1-3x^2}{(1+x^2)^3} \right]$$

$$\Rightarrow f'''(0) = -2$$

$$\therefore \tan^{-1}x = x - 2 \frac{x^3}{3!} + \dots$$

$$= x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

6. The power series expansion of  $\frac{\sin x}{x-\pi}$  about  $x=\pi$  is \_\_\_\_\_.

$$\text{Soln: } \text{Let, } x-\pi = t, \text{ then } f = \frac{\sin(\pi+t)}{t} \text{ about } t=0$$

$$= -\frac{\sin t}{t} \text{ about } t=0$$

$$= -\frac{1}{t} \left[ t - \frac{t^3}{3!} + \frac{t^5}{5!} - \frac{t^7}{7!} + \dots \right]$$

$$= -1 + \frac{t^2}{2!} - \frac{t^4}{4!} + \frac{t^6}{6!} - \dots$$

$$= -1 + \frac{(x-\pi)^2}{2!} - \frac{(x-\pi)^4}{4!} + \frac{(x-\pi)^6}{6!} - \dots$$

7. For the fun  $e^{-x}$ , linear approximation around  $x=2$  is \_\_\_\_\_

- (a)  $(3-x)e^{-2}$       (b)  $(1-x)e^{-2}$       (c)  $[\sqrt{3} + 2\sqrt{2} - (1+\sqrt{2})x]e^{-2}$

Sol<sup>n</sup>:→  $f(x) = \underbrace{f(a) + (x-a)f'(a)}_{\text{Linear approx.}} + \frac{(x-a)^2}{2!} f''(a) + \dots$

$$\therefore e^{-x} = f(2) + (x-2) \left. \frac{d}{dx} e^{-x} \right|_{x=2} = e^{-2} + (x-2)(-1)e^{-2} \\ = e^{-2}(3-x)$$

### 3. Definite Integrals:→

Theorem:→ Let  $f(x)$  is a continuous fun defined in  $[a, b]$  &  $F(x)$

be the anti-derivative of  $f(x)$  then  $\int_a^b f(x) dx = F(b) - F(a)$ .

Note:→  $\frac{d}{dx} \left[ \int_{u(x)}^{v(x)} f(x) dx \right] = f(v) \frac{dv}{dx} - f(u) \frac{du}{dx}$

### Properties of Definite Integrals:→

$$(i) \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$(ii) \text{ If } c \in (a, b) \text{ then, } \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

$$(iii) \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$(iv) \int_a^b \frac{f(x)}{f(x) + f(a+b-x)} dx = \frac{b-a}{2}$$

$$(v) \int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & ; \text{ if } f(x) \text{ is even} \\ 0 & , \text{ if } f(x) \text{ is odd} \end{cases}$$

$$(vi) \int_0^{2a} f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & ; \text{ if } f(2a-x) = f(x) \\ 0 & ; \text{ if } f(2a-x) = -f(x) \end{cases} \quad 12$$

$$(vii) \int_0^a x f(x) dx = \frac{a}{2} \int_0^a f(x) dx ; \text{ if } f(a-x) = f(x)$$

$$(viii) \int_0^{na} f(x) dx = n \int_0^a f(x) dx ; \text{ if } f(x+a) = f(x)$$

$$(ix) \int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx = \left[ \frac{n-1}{n} \times \frac{n-3}{n-2} \dots \times \frac{2}{3} (\text{or}) \frac{1}{2} \right] K$$

where,  $K = \begin{cases} 1 & ; \text{ if } n \text{ is odd} \\ \pi/2 & ; \text{ if } n \text{ is even} \end{cases}$

$$(x) \int_0^{\pi/2} \sin^m x \cos^n x dx = \left\{ \frac{[(m-1)(m-3) \dots 2(\text{or}) 1][(n-1)(n-3) \dots 2(\text{or}) 1]}{[(m+n)(m+n-2) \dots 2(\text{or}) 1]} \right\}$$

where,  $K = \begin{cases} \pi/2 & ; \text{ when } m \& n \text{ are even} \\ 0 & ; \text{ otherwise} \end{cases}$

Questions: →

$$1. \int_0^{\pi/2} \frac{\tan x}{\tan x + \cot x} dx = \underline{\hspace{2cm}}$$

Soln: → let  $f(x) = \tan x \Rightarrow f(0 + \frac{\pi}{2} - x) = \cot x$

$$2. I = \int_0^{\pi/2} \frac{f(x)}{f(x) + f(0 + \frac{\pi}{2} - x)} = \frac{\frac{\pi}{2} - 0}{2} = \frac{\pi}{4}.$$

2.  $\int_0^{\pi/2} \frac{dx}{1 + \sqrt{\cot x}}$  is \_\_\_\_\_.

Sol<sup>n</sup>:→  $I = \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx = \frac{\frac{\pi}{2} - 0}{2} = \frac{\pi}{4}$

3.  $\int_2^3 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{5-x}} dx$  is \_\_\_\_\_.

Sol<sup>n</sup>:→  $I = \frac{3-2}{2} = \frac{1}{2}$

4.  $\int_0^{\pi} |\cos x| dx$  is \_\_\_\_\_.

- (a) 1      (b) 0      (c) 2      (d) None.

Sol<sup>n</sup>:→  $I = \int_0^{\pi/2} \cos x dx + \int_{\pi/2}^{\pi} -\cos x dx$   
 $= \sin x \Big|_0^{\pi/2} - \sin x \Big|_{\pi/2}^{\pi}$   
 $= 1 - (-1) = 2$

5.  $\int_0^4 (|x| + |3-x|) dx$  is \_\_\_\_\_.

Sol<sup>n</sup>:→  $I = \int_0^4 x dx + \int_0^3 (3-x) dx + \int_3^4 (x-3) dx$   
 $= \frac{x^2}{2} \Big|_0^4 + \left[ 3x - \frac{x^2}{2} \right]_0^3 + \left[ \frac{x^2}{2} - 3x \right]_3^4$   
 $= 8 + \left[ 9 - \frac{9}{2} \right] + \left[ 8 - 12 - \frac{9}{2} + 9 \right]$   
 $= 34 - 21$

= 13

13

6.  $\int_0^n [x] dx$  is \_\_\_\_.

- (a)  $\frac{n(n+1)}{2}$       (b)  $\frac{n(n-1)}{2}$       (c)  $\frac{n}{2}$       (d) None.

Sol<sup>n</sup>:  $I = \int_0^1 0 \cdot dx + \int_1^2 1 \cdot dx + \int_2^3 2 \cdot dx + \dots + \int_{n-1}^n (n-1) \cdot dx$

$$= 0 + [2-1] + 2[3-2] + \dots + (n-1)[n-n+1]$$

$$= 1 + 2 + \dots + (n-1)$$

$$= \frac{n(n-1)}{2}$$

7.  $\int_0^1 x(1-x)^5 dx$  is \_\_\_\_.

- (a)  $\frac{1}{42}$       (b)  $\frac{1}{30}$       (c)  $\frac{1}{24}$       (d) None

Sol<sup>n</sup>:  $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

$$\int_0^1 x(1-x)^5 dx = \int_0^1 (1-x)x^5 dx = \frac{x^6}{6} \Big|_0^1 - \frac{x^7}{7} \Big|_0^1 = \frac{1}{42}$$

8.  $\int_0^{\pi/2} \log(\tan x) dx$  is \_\_\_\_.

Sol<sup>n</sup>: Replace  $x$  by  $(\frac{\pi}{2}-x)$ .

$$I = \int_0^{\pi/2} \log \cot x$$

$$2I = \int_0^{\pi/2} [\log \tan x + \log \cot x] dx = \int_0^{\pi/2} \log [\tan x \cdot \cot x] dx = 0$$

$$\Rightarrow I = 0.$$

9.  $\int_0^{\pi/4} \log(1+\tan x) dx$  is \_\_\_\_\_.

- (a)  $\frac{\pi}{8} \log 2$     (b)  $\frac{\pi}{4} \log 2$     (c)  $\frac{\pi}{2} \log 2$     (d) None

Sol<sup>n</sup>:  $\rightarrow I = \int_0^{\pi/4} \log(1+\tan(\frac{\pi}{4}-x)) dx$

$$= \int_0^{\pi/4} \log\left(1 + \frac{1-\tan x}{1+\tan x}\right) dx$$

$$= \int_0^{\pi/4} \log\left(\frac{2}{1+\tan x}\right) dx$$

$$= \int_0^{\pi/4} [\log 2 - \log(1+\tan x)] dx = \int_0^{\pi/4} \log 2 dx - \underbrace{\int_0^{\pi/4} \log(1+\tan x) dx}_I$$

$$\Rightarrow 2I = \int_0^{\pi/4} \log 2 dx = \frac{\pi}{4} \log 2$$

$$\Rightarrow I = \frac{\pi}{8} \log 2.$$

10.  $\int_{-1}^1 \frac{|x|}{x} dx$  is \_\_\_\_\_.

Sol<sup>n</sup>:  $\rightarrow I = 0$  (odd fun)

11.  $\int_{-\pi}^{\pi} \log \frac{(1+\sin x)}{(1-\sin x)} dx$  is \_\_\_\_\_.

Sol<sup>n</sup>:  $\rightarrow f(-x) = \log \frac{(1-\sin x)}{(1+\sin x)} = -\log \frac{(1+\sin x)}{(1-\sin x)} = -f(x) \Rightarrow$  odd fun

$\therefore I = 0$ .

12.  $\int_0^{\pi} \frac{x \sin x}{1+\cos^2 x} dx$  is \_\_\_\_\_.

Sol<sup>n</sup>:  $\rightarrow \int_0^a x f(x) dx = \frac{a}{2} \int_0^a f(x) dx$ ; if  $f(a-x) = f(x)$

$$f(\pi-x) = \frac{\sin x}{1+\cos^2 x} = f(x)$$

$$\therefore I = \frac{\pi}{2} \int_0^{\pi} \frac{\sin x}{1+\cos^2 x} dx$$

Put,  $\cos x = t \Rightarrow -\sin x dx = dt$

$$\Rightarrow I = \frac{\pi}{2} \int_{-1}^1 \frac{-dt}{1+t^2} = -\frac{\pi}{2} [\tan^{-1} t]_{-1}^1 = -\frac{\pi}{2} [\tan^{-1}(-1) - \tan^{-1}(1)] \\ = -\frac{\pi}{2} \left[ -\frac{\pi}{4} - \frac{\pi}{4} \right] = \frac{\pi^2}{4}$$

13.  $\int_0^{\pi} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}$  is \_\_\_\_\_.

(a)  $\frac{\pi}{ab}$

(b)  $\pi ab$

(c)  $\frac{2\pi}{ab}$

(d) None

Sol:  $I = \int_0^{\pi} \frac{\sec^2 x dx}{a^2 + b^2 \tan^2 x}$

$$\int_0^{2a} f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & ; \text{ if } f(2a-x) = f(x) \\ 0 & ; \text{ if } f(2a-x) = -f(x) \end{cases}$$

~~f(x)~~  $f(\pi-x) = f(x)$

$$\therefore I = 2 \int_0^{\pi/2} \frac{\sec^2 x}{a^2 + b^2 \tan^2 x} dx$$

Let,  $\tan x = t$

$$\sec^2 x dx = dt$$

$$\therefore I = 2 \int_0^{\infty} \frac{dt}{a^2 + (bt)^2} = 2 \times \frac{1}{a} \left[ \frac{\tan^{-1}(bt/a)}{b} \right]_0^{\infty} = \frac{2}{ab} \left[ \frac{\pi}{2} - 0 \right] = \frac{\pi}{ab}$$

14. If  $f(t)$  is a continuous fun defined in  $[0, 1]$  then  $\lim_{t \rightarrow 0} \frac{1}{t} \int_0^t f(t) dt$

- (a) 0    (b)  $\infty$     (c)  $f(0)$     (d)  $f'(1)$

$$\text{Soln: } \lim_{t \rightarrow 0} \frac{\int_0^t f(t) dt}{t} = \lim_{t \rightarrow 0} \frac{\frac{d}{dt} \left\{ \int_0^t f(t) dt \right\}}{1}$$

$$= \lim_{t \rightarrow 0} [f(t) \times 1 - f(0) \times 0]$$

$$= f(0)$$

$$15. \int_0^{\pi/2} \sin^8 x dx$$

$$\text{Soln: } I = \frac{7}{8} \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} =$$

$$16. \int_0^{\pi/2} \cos^7 x dx$$

$$\text{Soln: } I = \frac{6}{7} \times \frac{4}{5} \times \frac{2}{3} \times 1 =$$

$$17. \int_0^{\pi/2} \sin^5 x \cos^9 x dx$$

$$\text{Soln: } I = \frac{(4 \times 2) \times (8 \times 6 \times 4 \times 2)}{14 \times 12 \times 10 \times 8 \times 6 \times 4 \times 2} \times 1 =$$

$$18. \int_0^{\pi/2} \sin^6 x \cos^3 x dx$$

$$\text{Soln: } I = \frac{(5 \times 3 \times 1) \times (2)}{9 \times 7 \times 5 \times 3 \times 1} \times 1 =$$

$$19. \int_0^{\pi/2} \sin^5 x \cos^8 x dx$$

$$\text{Soln: } I = \frac{(4 \times 2) \times (7 \times 5 \times 3 \times 1)}{13 \times 11 \times 9 \times 7 \times 5 \times 3 \times 1} \times 1 =$$

$$20. \int_0^{\pi/2} \sin^6 x \cdot \cos^8 x \, dx = \underline{\hspace{2cm}}$$

15

$$\text{Soln: } \rightarrow I = \frac{(5 \times 3 \times 1) (7 \times 5 \times 3 \times 1)}{14 \times 12 \times 10 \times 8 \times 6 \times 4 \times 2} \times \frac{\pi}{2} =$$

$$21. \int_{-\pi}^{\pi} \sin^4 x \, dx = \underline{\hspace{2cm}}$$

$$\text{Soln: } \rightarrow I = 2 \int_0^{\pi} \sin^4 x \, dx$$

$$\int_0^{2a} f(x) \, dx = \begin{cases} 2 \int_0^a f(x) \, dx & ; \text{ if } f(2a-x) = f(x) \\ & ; \text{ if } f(2a-x) = -f(x) \end{cases}$$

$$\therefore \sin^4(\pi-x) = \sin^4 x$$

$$\therefore I = 2 \times 2 \times \int_0^{\pi/2} \sin^4 x \, dx = 4 \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} =$$

$$22. \int_0^{\pi} \sin^4 x \cos^3 x \, dx = \underline{\hspace{2cm}}$$

$$\text{Soln: } \rightarrow f(\pi-x) = \sin^4 x (-\cos x)^3 = -f(x)$$

$$\therefore I = 0.$$

$$23. \int_0^{\pi} \sin^3 x \cos^4 x \, dx = \underline{\hspace{2cm}}$$

$$\text{Soln: } \rightarrow f(\pi-x) = \sin^3 x \cos^4 x = f(x)$$

$$\therefore I = 2 \int_0^{\pi/2} \sin^3 x \cos^4 x \, dx = 2 \times \frac{(2 \times 3 \times 1)}{7 \times 5 \times 3 \times 1} \times 1 =$$

$$24. \int_{-\frac{2\pi}{2}}^{\frac{2\pi}{2}} \sin^4 x \cos^8 x \, dx$$

$$\text{Solt:} \rightarrow I = 2 \int_0^{\frac{\pi}{2}} \sin^4 x \cos^8 x \, dx = 2 \times 2 \times \int_0^{\frac{\pi}{2}} \sin^4 x \cos^8 x \, dx \\ = 2 \times 2 \times 2 \times \int_0^{\frac{\pi}{2}} \sin^4 x \cos^8 x \, dx \\ = 8 \times \frac{(3 \times 1) \times (7 \times 5 \times 3 \times 1)}{12 \times 10 \times 8 \times 6 \times 4 \times 2} \times \frac{\pi}{2}$$

$$\text{Note:} \rightarrow \int_a^b \sin^4 x \cos^8 x \, dx = K \int_0^{\frac{\pi}{2}} \sin^4 x \cos^8 x \, dx$$

$$\text{where, } K = \frac{b-a}{\frac{\pi}{2}}$$

$$25. \int_0^{\frac{\pi}{2}} \sin x \cos^3 x \, dx$$

solt: Let  $\sin x = t \Rightarrow \cos x \, dx = dt$

$$\therefore I = \int_0^1 \sqrt{t} (1-t^2) \, dt = \frac{t^{3/2}}{3/2} \Big|_0^1 - \frac{t^3}{3} \Big|_0^1 = \frac{2}{3} - \frac{2}{7} =$$

### Improper Integrals:

First Kind:  $\int_a^b f(x) \, dx$  if  $a = -\infty$  (or)  $b = \infty$  (or) both

$$\text{i.e. } \int_{-\infty}^b f(x) \, dx, \int_a^{\infty} f(x) \, dx, \int_{-\infty}^{\infty} f(x) \, dx$$

Second Kind:  $\rightarrow \int_a^b f(x) dx$  if  $a \leq b$  are finite but  $f(x)$  is infinite for some  $x \in [a, b]$ . 10

$$\text{Ex: } \int_{-1}^1 \log(1+x) dx, \quad \int_0^1 \frac{1}{1-x} dx, \quad \int_0^3 \frac{1}{x^2 - 5x + 4} dx$$

Convergence of an Improper Integrals:  $\rightarrow$

(i) If  $\int_a^b f(x) dx = \text{finite}$ , then, it is a convergent improper integral.

(ii) If  $\int_a^b f(x) dx = \text{infinite}$ , then, it is a divergent improper integral.

Questions:  $\rightarrow$

1. Find the convergence of following improper integrals

$$(i) \int_0^\infty \frac{1}{a^2 + x^2} dx$$

$$\text{Soln: } I = \frac{1}{a} \tan^{-1} \frac{x}{a} \Big|_0^\infty = \frac{1}{a} \left[ \frac{\pi}{2} - 0 \right] = \frac{\pi}{2a} = \text{finite}$$

which is convergent.

$$(ii) \int_0^\infty x \sin x dx$$

$$\text{Soln: } I = x \left[ (-\cos x) - (1)(-\sin x) \right] \Big|_0^\infty = \text{infinite}$$

which is divergent improper integral.

$$(iii) \int_{-\infty}^0 e^{ax} \cos px dx = \underline{\hspace{2cm}}$$

$$\text{Soln: } \rightarrow \int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2+b^2} [a \cos bx + b \sin bx]$$

$$\therefore I = \left[ \frac{e^{ax}}{a^2+p^2} [a \cos px + p \sin px] \right]_0^\infty \\ = \frac{a}{a^2+p^2} - 0 = \text{finite} \\ \text{i.e. convergence}$$

$$(iv) \int_{-1}^1 \sqrt{\frac{1+x}{1-x}} dx$$

$$\text{Soln: } \rightarrow I = \int_{-1}^1 \frac{1+x}{\sqrt{1-x^2}} dx = \int_{-1}^1 \frac{1}{\sqrt{1-x^2}} dx + \int_{-1}^1 \frac{x}{\sqrt{1-x^2}} dx \\ = 2 \int_0^1 \frac{1}{\sqrt{1-x^2}} dx + 0 \\ = 2 \left[ \sin^{-1} x \right]_0^1 = 2 \cdot \frac{\pi}{2} = \pi = \text{finite}$$

i.e. convergent improper integral.

$$(v) \int_{-1}^1 \frac{1}{x^2} dx = \underline{\hspace{2cm}}$$

$$\text{Soln: } \rightarrow I = \int_{-1}^0 \frac{1}{x^2} dx + \int_0^1 \frac{1}{x^2} dx = \left[ -\frac{1}{x} \right]_{-1}^0 + \left[ -\frac{1}{x} \right]_0^1 = \text{infinite}$$

i.e. divergent improper integral

12

$$(vi) \int_0^3 \frac{1}{x^2 - 3x + 2} dx$$

$$\text{Soln: } \rightarrow I = \int_0^3 \frac{1}{(x-1)(x-2)} dx = \int_0^1 \frac{dx}{(x-1)(x-2)} + \int_1^2 \frac{dx}{(x-1)(x-2)} + \int_2^3 \frac{dx}{(x-1)(x-2)}$$

$$\int \frac{dx}{(x-1)(x-2)} = \int \frac{dx}{x-2} - \int \frac{dx}{x-1} = \log \left( \frac{x-2}{x-1} \right)$$

$$\therefore I = \left. \log \left( \frac{x-2}{x-1} \right) \right|_0^1 + \left. \log \left( \frac{x-2}{x-1} \right) \right|_1^2 + \left. \log \left( \frac{x-2}{x-1} \right) \right|_2^3 = \text{infinite}$$

i.e., divergent improper integral.

### Comparison Test:

for first kind of improper integrals:

(a) Let  $0 \leq f(x) \leq g(x)$ , then,

(i)  $\int_a^b f(x) dx$  converges if  $\int_a^b g(x) dx$  is convergent

(ii)  $\int_a^b g(x) dx$  diverges if  $\int_a^b f(x) dx$  is divergent

(b) Limit form: → let  $f(x)$  &  $g(x)$  be two positive fun s.t.

$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = l$  (non-zero, finite) then  $\int_a^b f(x) dx$  &

$\int_a^b g(x) dx$  both convergent or divergent together.

Questions:

$$1. \int_1^\infty e^{-x^2} dx =$$

$$\text{Soln: } \rightarrow e^{-x^2} \geq e^{-x} \quad \forall x \geq 1$$

$$\Rightarrow e^{-x^2} \leq e^{-x} \quad \forall x \geq 1$$

$$\int_1^\infty e^{-x} dx = \left[ -e^{-x} \right]_1^\infty = 1 \Rightarrow \int_1^\infty e^{-x} dx \text{ is convergent}$$

$\therefore \int_1^\infty e^{-x^2} dx$  is also convergent.

$$2. \int_2^\infty \frac{1}{\log x} dx$$

$$\text{Soln: } \log x < \infty \quad \forall x \geq 2$$

$$\Rightarrow \frac{1}{\log x} > \frac{1}{x} \quad \forall x \geq 2$$

$$\int_2^\infty \frac{1}{x} dx = \left[ \log x \right]_2^\infty = \infty \Rightarrow \text{divergent}$$

$\therefore \int_2^\infty \frac{dx}{\log x}$  is also divergent.

$$3. \int_1^\infty \frac{1}{x^2(e^{-x}+1)} dx$$

$$\text{Soln: } x^2(e^{-x}+1) > x^2(0+1) \quad \forall x > 1$$

$$\Rightarrow \frac{1}{x^2(e^{-x}+1)} < \frac{1}{x^2} \quad \forall x > 1$$

} Method- I

$$\text{Method-II: Let } g(x) = x^2, \quad \frac{f(x)}{g(x)} = \frac{1}{e^{-x}+1}$$

so that,  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$  thus,  $g(x)$  will gives the nature of  $f(x)$ .

$$\therefore \int_1^\infty \frac{1}{x^2} dx = \left[ -\frac{1}{x} \right]_1^\infty = 1 \Rightarrow \text{convergent}$$

Hence,  $f(x)$  is also convergent.

$$Q. 4. \int_1^{\infty} \frac{x \tan^{-1} x}{\sqrt{4+x^3}} dx$$

18

$$\text{Soln: } \rightarrow f(x) = \frac{x \tan^{-1} x}{x \sqrt{x} \sqrt{\frac{4}{x^3} + 1}} = \frac{\tan^{-1} x}{\sqrt{x} \sqrt{\frac{4}{x^3} + 1}}$$

$$\text{let } g(x) = \frac{1}{\sqrt{x}}$$

$$\frac{f(x)}{g(x)} = \frac{\tan^{-1} x}{\sqrt{\frac{4}{x^3} + 1}} \Rightarrow \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \pi/2$$

$$\int_1^{\infty} \frac{1}{\sqrt{x}} dx = 2\sqrt{x} \Big|_1^{\infty} = \text{infinite} \Rightarrow \text{divergent}$$

$\therefore f(x)$  is also divergent

Comparison Test for the second kind of improper integral : →

(b) Limit form: → Let  $f(x)$  &  $g(x)$  be two +ve fun s.t.

(i) 'a' is a point of discontinuity and  $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = l_1$

(non-zero & finite)

(ii) 'b' is a point of discontinuity and  $\lim_{x \rightarrow b^-} \frac{f(x)}{g(x)} = l_2$

(non-zero & finite)

then,  $\int_a^b f(x) dx$  &  $\int_a^b g(x) dx$  both converge (or) diverge together.

Questions: →

$$1. \int_0^{\pi/2} \frac{\sin x}{x\sqrt{x}} dx$$

Soln: →  $\frac{\sin x}{x} \leq 1 \quad \forall x > 0$

$$\Rightarrow \frac{\sin x}{x\sqrt{x}} \leq \frac{1}{\sqrt{x}} \quad \forall x > 0$$

$$\int_0^{\pi/2} \frac{1}{\sqrt{x}} dx = 2\sqrt{x} \Big|_0^{\pi/2} = 2\sqrt{\frac{\pi}{2}} = \text{finite} \Rightarrow \text{convergent}$$

∴  $\int_0^{\pi/2} \frac{\sin x}{x\sqrt{x}} dx$  is also convergent

$$2. \int_1^2 \frac{\sqrt{x}}{\log x} dx$$

Soln: →  $\frac{1}{\log x} > \frac{1}{x} \quad \forall x > 1$

$$\Rightarrow \frac{\sqrt{x}}{\log x} > \frac{1}{\sqrt{x}} \quad \forall x > 1$$

$$\int_1^2 \frac{1}{\sqrt{x}} dx = 2\sqrt{x} \Big|_1^2 = 2\sqrt{2} - 2 = \text{finite} \Rightarrow \text{convergent}$$

∴  $\int_1^2 \frac{\sqrt{x}}{\log x} dx$  may/may not be convergent.

Thus, first method fails.

$$\lim_{x \rightarrow 1^+} \frac{f(x)}{g(x)} = 1$$

Let,  $g(x) = \frac{1}{x \log x}$

$$\Rightarrow \frac{f(x)}{g(x)} = x \sqrt{x}$$

$$\int_1^2 \frac{1}{x \log x} dx ; \text{ Put } \log x = t$$

$$\Rightarrow \frac{1}{x} dx = dt$$

$$\int_1^2 \frac{1}{x \log x} dx = \int_0^{\log 2} \frac{1}{t} dt = \log t \Big|_0^{\log 2} = \text{infinite} \Rightarrow \text{divergent}$$

$$\therefore \int_1^2 \frac{\sqrt{x}}{\log x} dx \text{ is also divergent}$$

3. Which of the following fun is strictly bounded :-

- (a)  $x^2$  (b)  $e^x$  (c)  $\frac{1}{x}$  (d)  $e^{-x^2}$

4. Which of the following integrals is unbounded :-

$$(a) \int_0^{\pi/4} \tan x dx \quad (b) \int_0^{\infty} \frac{1}{1+x^2} dx \quad (\text{c}) \int_0^1 \frac{1}{1-x} dx \quad (\text{d}) \int_0^{\infty} x e^{-x} dx$$

$$\text{Solt: } \rightarrow (a) I = \log \sec x \Big|_0^{\pi/4} = \log \sqrt{2}$$

$$(b) I = \tan^{-1} x \Big|_0^{\infty} = \pi/2$$

$$(c) I = -\log(1-x) \Big|_0^1 = \text{infinite}$$

$$(d) I = x e^{-x} \Big|_{-1}^{\infty} - \left( 1 \right) \frac{e^{-x}}{(-1)^2} \Big|_0^{\infty} = 1$$

$$5. \text{ Consider the integrals } I_1 = \int_1^{\infty} \frac{1}{x^2(e^x+1)} dx \quad \& \quad I_2 = \int_1^{\infty} \frac{x+1}{x \sqrt{x}} dx$$

then which of the following is true :-

(a)  $I_1$  &  $I_2$  are convergent ~~(b)  $I_1$  &  $I_2$  are divergent~~

~~(b)  $I_1$  is convergent,  $I_2$  is divergent~~

(c)  $I_1$  is divergent,  $I_2$  is convergent

(d)  $I_1$  &  $I_2$  are divergent

$$\text{Soln: } x^2(e^x+1) > x^2(0+1) \quad \forall x \geq 1$$

$$\Rightarrow \frac{1}{x^2(e^x+1)} < \frac{1}{x^2} \quad \forall x \geq 1$$

$$\int_1^\infty \frac{1}{x^2} dx = -\frac{1}{x} \Big|_1^\infty = 1 \Rightarrow \text{convergent}$$

$\therefore I_1$  is convergent.

$$I_2 = \int_1^\infty \frac{x}{x\sqrt{x}} dx + \int_1^\infty \frac{1}{x\sqrt{x}} dx = \int_1^\infty \frac{1}{\sqrt{x}} dx + \int_1^\infty \frac{1}{x\sqrt{x}} dx$$

$$\downarrow$$

$$2\sqrt{x} \Big|_1^\infty \Rightarrow \text{Divergent}$$

$\therefore I_2$  is divergent.

6. consider,  $I_1 = \int_0^1 \frac{1}{x^{1/3}} dx, I_2 = \int_1^\infty \frac{1}{x} dx, I_3 = \int_0^1 x \log x dx$

then which of the following is convergent:-

(a)  $I_1 \& I_2$

(b)  $I_2 \& I_3$

(c)  $I_1 \& I_3$

(d) Only  $I_1$

$$\text{Soln: } I_1 = \left. \frac{x^{2/3}}{2/3} \right|_0^1 = \text{finite}$$

$$I_2 = \int_0^\infty \frac{1}{x} dx + \int_1^\infty \frac{1}{x} dx = \log x \Big|_1^\infty + \log x \Big|_0^\infty = \text{infinite}$$

$$I_3 = \left. \log x \cdot \frac{x^2}{2} \right|_0^1 - \int_0^1 \frac{1}{x} \cdot \frac{x^2}{2} dx = \left[ \frac{x^2}{2} \log x - \frac{x^2}{4} \right]_0^1$$

$$= -\frac{1}{4} - \left[ \lim_{x \rightarrow 0} \frac{x^2}{2} (\log x) \right]$$

$$\lim_{x \rightarrow 0} \frac{\log x}{2/x^2} = \lim_{x \rightarrow 0} \frac{1/x}{-4/x^3} = \lim_{x \rightarrow 0} -\frac{x^2}{4} = 0 \quad 20$$

$\therefore I_3 = \text{finite}$

### Gamma Function:

$$\Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx \quad (n > 0)$$

Note: (i)  $\Gamma(1) = 1$

$$(ii) \Gamma(\frac{1}{2}) = \sqrt{\pi}$$

$$(iii) \Gamma(n+1) = n\Gamma(n) \quad \forall n > 0$$

$$(iv) \Gamma(n+1) = n! \quad \forall n \in \mathbb{Z}^+$$

$$(v) \int_0^\infty e^{-ax} x^{n-1} dx = \frac{\Gamma(n)}{a^n}$$

### Questions:

$$1. \int_0^\infty e^{-x^2} dx = \underline{\hspace{2cm}}$$

$$\text{Soln:} \rightarrow \text{Let } x^2 = t \Rightarrow 2x dx = dt \Rightarrow dx = \frac{1}{2} t^{-\frac{1}{2}} dt$$

$$\therefore I = \int_0^\infty e^{-t} \cdot \frac{1}{2} t^{-\frac{1}{2}} dt = \frac{1}{2} \int_0^\infty e^{-t} t^{\frac{1}{2}-1} dt = \frac{1}{2} \Gamma(\frac{1}{2}) = \frac{\sqrt{\pi}}{2}$$

$$\text{Note:} \rightarrow \int_{-\infty}^\infty e^{-x^2} dx = 2 \int_0^\infty e^{-x^2} dx = \sqrt{\pi}$$

$$2. \int_0^\infty e^{-2x^2} x^7 dx = \underline{\hspace{2cm}}$$

$$\text{Soln:} \rightarrow \text{Let } 2x^2 = t \Rightarrow 4x dx = dt$$

$$\therefore I = \int_0^\infty e^{-t} \left(\frac{t}{2}\right)^3 \frac{dt}{4} = \frac{1}{32} \int_0^\infty e^{-t} t^3 dt = \frac{1}{32} \Gamma(4) = \frac{3!}{32} = \frac{6}{32}$$

$$3. \int_0^1 (x \log x)^4 dx = \underline{\hspace{2cm}}$$

Soln: Let,  $\log x = -t \Rightarrow x = e^{-t} \Rightarrow dx = -e^{-t} dt$

$$\therefore I = \int_{-\infty}^0 [e^{-t}(-t)]^4 (-e^{-t}) dt = \int_0^\infty e^{-5t} t^{5-1} dt = \frac{\Gamma(5)}{5^5} = \frac{4!}{5^5}$$

$$4. \int_0^\infty 5^{-4x^2} dx = \underline{\hspace{2cm}}$$

$$\text{Soln:} \text{ Let, } 5^{-4x^2} = e^{-t} \Rightarrow -4x^2 \log 5 = -t \Rightarrow x = \frac{1}{2\sqrt{\log 5}} \sqrt{t}$$

$$\Rightarrow dx = \frac{1}{2\sqrt{\log 5}} \frac{1}{2\sqrt{t}} dt$$

$$\begin{aligned} I &= \int_0^\infty e^{-t} \cdot \frac{1}{2\sqrt{\log 5}} \cdot \frac{t^{-\frac{1}{2}}}{2} dt = \frac{1}{4\sqrt{\log 5}} \int_0^\infty e^{-t} t^{\frac{1}{2}-1} dt \\ &= \frac{\sqrt{\pi}}{4\sqrt{\log 5}} \end{aligned}$$

$$\text{Note:} \log a^0 = \begin{cases} \infty &; a < 1 \\ -\infty &; a > 1 \end{cases}$$

Beta function:  $\rightarrow$

$$\beta(m, n) = \int_0^1 x^{m-1} (1-x)^{n-1} dx \quad (m > 0, n > 0)$$

$$\text{Note:} (i) \beta(m, n) = \beta(n, m)$$

$$(ii) \beta(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$$

$$(iii) \beta(m, n) = \int_0^\infty \frac{x^{m-1}}{(1+x)^{m+n}} dx = \int_0^\infty \frac{x^{n-1}}{(1+x)^{m+n}} dx$$

$$(iv) \beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1}\theta \cos^{2n-1}\theta d\theta$$

21

$$\text{i.e. } \int_0^{\pi/2} \sin^p \theta \cos^q \theta d\theta = \frac{1}{2} \beta\left(\frac{p+1}{2}, \frac{q+1}{2}\right)$$

$$(p > -\frac{1}{2}, q > -\frac{1}{2})$$

Questions:

$$1. \int_0^2 x^7 (16-x^4)^{10} dx = \underline{\hspace{2cm}}$$

$$\text{Soln:} \rightarrow \text{Let, } x^4 = 16t \Rightarrow 4x^3 dx = 16 dt \Rightarrow x^3 dx = 4 dt$$

$$1 = \int_0^1 16t (16-16t)^{10} 4 dt = 16 \times 4 \int_0^1 t (1-t)^{10} dt$$

$$= 4 \times 16^{11} \times \beta(2, 11)$$

$$= 4 \times 16^{11} \times \frac{12!}{13!}$$

$$= 4 \times 16^{11} \times \frac{1 \times 10!}{12!}$$

$$2. \int_0^\infty \frac{x^3 (1+x^5)}{(1+x)^{13}} dx = \underline{\hspace{2cm}}$$

$$\text{Soln:} \rightarrow \int_0^\infty \frac{x^3}{(1+x)^{13}} dx + \int_0^\infty \frac{x^8}{(1+x)^{13}} dx = \int_0^\infty \frac{x^{4-1}}{(1+x)^{4+9}} dx + \int_0^\infty \frac{x^{9-1}}{(1+x)^{9+4}} dx$$

$$= \beta(4, 9) + \beta(9, 4)$$

$$= 2 \beta(4, 9) = 2 \times \frac{14 \times 19}{13} = 2 \times \frac{3! \times 8!}{12!}$$

$$3. \int_0^{\infty} \left( \frac{x}{1+x^2} \right)^3 dx = \dots$$

Soln: → Let  $x = \tan \theta \Rightarrow dx = \sec^2 \theta d\theta$

$$\begin{aligned} \therefore I &= \int_0^{\pi/2} \left( \frac{\tan \theta}{\sec^2 \theta} \right)^3 \sec^2 \theta d\theta = \int_0^{\pi/2} (\tan^3 \theta / \sec^4 \theta) d\theta \\ &= \int_0^{\pi/2} \sin^3 \theta \cos \theta d\theta \\ &= \frac{1}{2} \beta \left( \frac{3+1}{2}, \frac{1+1}{2} \right) = \frac{1}{2} \beta (2, 1) \\ &= \frac{1}{2} \times \frac{\sqrt{2} \sqrt{1}}{\sqrt{3}} = \frac{1}{2} \times \frac{1 \times 1}{2!} = \frac{1}{4} \end{aligned}$$

#### 4. Partial Differentiation: →

Let  $z = f(x, y)$  then

$$z_x = \frac{\partial z}{\partial x} = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

$$z_y = \frac{\partial z}{\partial y} = \lim_{k \rightarrow 0} \frac{f(x, y+k) - f(x, y)}{k}$$

Similarly,  $\frac{\partial^2 z}{\partial x^2}$ ,  $\frac{\partial^2 z}{\partial y^2}$ ,  $\frac{\partial^2 z}{\partial x \partial y}$  and so on.

#### Homogeneous Function: →

$$\text{Ex:- (i) } x^2 + xy + y^2 ; n = 2$$

$$(ii) 2x^3 + xy^2 + z^3 ; n = 3$$

$$(iii) \frac{xy^2 - y^3}{2x + 3y} ; n = 3-1 = 2$$

If  $f(kx, ky) = k^n f(x, y)$  then  $f(x, y)$  is a homogeneous function with degree 'n'.

Note → (i) If  $f(x, y)$  is a homogeneous function with degree 'n', then,

$$f(x, y) = \begin{cases} x^n \phi(y/x) \\ y^n \psi(x/y) \end{cases}$$

Euler's Theorem → If  $f(x, y)$  is a homogeneous function with degree 'n', then,

$$(a) x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} = n f$$

$$(b) x^2 \frac{\partial^2 f}{\partial x^2} + 2xy \frac{\partial^2 f}{\partial x \partial y} + y^2 \frac{\partial^2 f}{\partial y^2} = n(n-1)f$$

Note → If  $u(x, y) = f(x, y) + g(x, y) + h(x, y)$ , where,  $f, g$  &  $h$

are homogeneous functions with degree  $m, n$  &  $p$  respectively, then,

$$(a) x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = mf + ng + ph$$

$$(b) x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = m(m-1)f + n(n-1)g + p(p-1)h$$

Note → If  $f(u)$  is a homogeneous function in two variables  $x$  &  $y$  with degree 'n', then,

$$(a) x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} = F(u)$$

$$(b) x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = F(u) [F'(u) - 1]$$

Total Differentiation → If  $z = f(x, y)$  where  $x = \phi(t)$ ,  $y = \psi(t)$ ,

then, the total derivative of 'z' w.r.t. 't' is

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dt}$$

Total differential coefficient,

$$dz = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

Note: → (i) If  $f(x, y) = c$  is an implicit fun, then,

$$\frac{dy}{dx} = -\frac{f_x}{f_y}$$

(ii) If  $z = f(x, y)$ , where  $x = \phi(u, v)$  &  $y = \psi(u, v)$ , then,

$$\frac{\partial z}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u}$$

$$\text{&} \quad \frac{\partial z}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v}$$

Questions: →

1. If  $z = e^x \sin y$ , where  $x = \log t$  &  $y = t^2$ , then,  $\frac{dz}{dt} = ?$

Soln: →  $\frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \cdot \frac{dy}{dt}$

$$= e^x \sin y \times \frac{1}{t} + e^x \cos y \times 2t$$

$$= \frac{e^x \sin y}{t} + 2te^x \cos y$$

$$= \frac{e^x}{t} (\sin y + 2t^2 \cos y)$$

$$= \sin y + 2y \cos y \quad \left\{ \because t = e^x \text{ & } t^2 = y \right\}$$

2. The total derivative of  $x^2 y$  wrt.  $x$ , where,  $x$  &  $y$  are connected by the relation  $x^2 + xy + y^2 = 1$  is \_\_\_\_\_.

Soln: → Let,  $u = x^2 y$

$$\frac{du}{dx} = \frac{\partial u}{\partial x} \cdot \frac{dx}{dx} + \frac{\partial u}{\partial y} \cdot \frac{dy}{dx}$$

$$= 2xy + x^2 \frac{dy}{dx}$$

Now,  $f(x, y) = x^2 + xy + y^2 = 1$

23

$$\therefore \frac{dy}{dx} = -\frac{fx}{fy} = -\frac{2x+y}{x+2y}$$

$$\therefore \frac{du}{dx} = 2xy + x^2 \left( -\frac{2x+y}{x+2y} \right) = 2xy - x^2 \left( \frac{2x+y}{x+2y} \right).$$

3. If  $u = f(x+cy) + g(x-cy)$ , then,  $\frac{u_{xx}}{u_{yy}} = \underline{\hspace{2cm}}$

- (a)  $c^{-2}$  (b)  $c^2$  (c)  $-c^{-2}$  (d)  $-c^2$

Soln: → Let,  $r = x+cy$ ,  $s = x-cy$

$$u = f(r) + g(s)$$

$$u_x = f'(r) \frac{\partial r}{\partial x} + g'(s) \frac{\partial s}{\partial x} = f'(r) + g'(s)$$

$$u_{xx} = f''(r) + g''(s)$$

$$u_y = f'(r) \frac{\partial r}{\partial y} + g'(s) \frac{\partial s}{\partial y} = c f'(r) - c g'(s)$$

$$u_{yy} = c^2 f''(r) + c^2 g''(s)$$

$$\therefore \frac{u_{xx}}{u_{yy}} = \frac{1}{c^2} = c^{-2}$$

4. If  $u = f(2x-3y, 3y-4z, 4z-2x)$ , then,  $6u_x + 4u_y = \underline{\hspace{2cm}}$

- (a)  $3u_z$  (b)  $4u_z$  (c)  $-3u_z$  (d)  $-4u_z$

Soln: →  $r = 2x-3y$ ,  $s = 3y-4z$ ,  $t = 4z-2x$

$$u = f(r, s, t)$$

$$\begin{aligned} u_x &= \frac{\partial f}{\partial r} \cdot \frac{\partial r}{\partial x} + \frac{\partial f}{\partial s} \cdot \frac{\partial s}{\partial x} + \frac{\partial f}{\partial t} \cdot \frac{\partial t}{\partial x} \\ &= 12f_r(2) + f_s(0) + 12f_t(-2) \end{aligned}$$

$$\therefore 6u_x = 12f_r - 12f_t$$

$$u_y = \frac{\partial f}{\partial r} \cdot \frac{\partial r}{\partial y} + \frac{\partial f}{\partial s} \cdot \frac{\partial s}{\partial y} + \frac{\partial f}{\partial t} \cdot \frac{\partial t}{\partial y}$$

$$= f_r(-3) + f_s(3) + f_t(0)$$

$$\therefore 4u_y = -12f_r + 12f_s$$

$$\therefore 6u_x + 4u_y = 12f_s - 12f_t$$

$$u_z = \frac{\partial f}{\partial r} \cdot \frac{\partial r}{\partial z} + \frac{\partial f}{\partial s} \cdot \frac{\partial s}{\partial z} + \frac{\partial f}{\partial t} \cdot \frac{\partial t}{\partial z}$$

$$= f_r(0) + f_s(-4) + f_t(4)$$

$$= -4f_s + 4f_t$$

5. If  $u = \frac{y}{z} + \frac{z}{x}$ , then,  $xu_x + yu_y + zu_z = \underline{\quad}$

- (a)  $\frac{xy}{z^2}$     (b)  $\frac{yz}{x^2}$     (c)  $\frac{xz}{y^2}$     (d)  $\checkmark 0$

Soln:  $\rightarrow xu_x + yu_y + zu_z = 0 \cdot u = 0 \quad (\because n=0)$

6. If  $\mu = \frac{x^2y}{x^{5/2} + y^{5/2}}$ , then,  $x^2\mu_{xx} + 2xy\mu_{xy} + y^2\mu_{yy} = \underline{\quad}$

- (a)  $\frac{3}{4}\mu$     (b)  $\checkmark -\frac{1}{4}\mu$     (c)  $-\frac{3}{4}\mu$     (d)  $\frac{1}{4}\mu$

Soln:  $\rightarrow n = 3 - \frac{5}{2} = \frac{1}{2}$

$$\therefore x^2\mu_{xx} + 2xy\mu_{xy} + y^2\mu_{yy} = n(n-1)\mu = \frac{-1}{4}\mu.$$

7. If  $u = \operatorname{cosec}^{-1} \left[ \frac{x^{1/4} - y^{1/4}}{x^{1/5} + y^{1/5}} \right]$ , then,  $xu_x + yu_y = \underline{\quad}$

- (a)  $-\frac{1}{20}u$     (b)  $-\frac{1}{20}\cot u$     (c)  $\frac{1}{20}\tan u$     (d)  $\checkmark -\frac{1}{20}\tan u$

Soln:  $\rightarrow \operatorname{cosec} u = \frac{x^{1/4} - y^{1/4}}{x^{1/5} + y^{1/5}}$

$$\Rightarrow n = \frac{1}{4} - \frac{1}{5} = \frac{1}{20}$$

$$\therefore x u_x + y u_y = n \frac{f(u)}{f'(u)} = \frac{1}{20} \times \frac{\csc u}{-\csc u \cot u} = -\frac{1}{20} \tan u.$$

24

8. If  $u = \log\left(\frac{x^2}{y}\right)$ , then,  $x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = \underline{\quad}$ .

- (a)  $u$     (b) 0    (c)  $-1$     (d) 1

Soln:  $\rightarrow e^u = \left(\frac{x^2}{y}\right) \Rightarrow n = 2-1 = 1$

$$x u_x + y u_y = n \frac{f(u)}{f'(u)} = \frac{e^u}{e^u} = 1 = f(u).$$

$$\therefore x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = F(u) [F'(u)-1] = -1.$$

9. If  $z = x^n f(y/x) + y^{-n} g(x/y)$ , then,  $x z_x + y z_y + x^2 z_{xx} + 2xy z_{xy} + y^2 z_{yy} = \underline{\quad}$ .

is  $\underline{\quad}$ .

- (a)  $n(n-1)z$     (b)  $n^2 z$     (c)  $n(n+1)z$     (d)  $nz$

Soln:  $\rightarrow$  The given fun is the sum of two homogeneous fun having degree  $n$  &  $-n$  respectively.

$$\therefore x z_x + y z_y + x^2 z_{xx} + 2xy z_{xy} + y^2 z_{yy} = (nf - ng) + [n(n-1)f] \\ + [(-n)(-n-1)g]$$

$$= n^2 z$$

10. If  $u = \sin^{-1}(xy) + \cos^{-1}(y/x)$ , then,  $\frac{u_x}{u_y} = \underline{\quad}$ .

- (a)  $-\frac{y}{x}$     (b)  $-\frac{x}{y}$     (c)  $\frac{y}{x}$     (d)  $\frac{x}{y}$

Soln:  $\rightarrow x u_x + y u_y = 0 \cdot u = 0 \quad (\because n=0)$

$$\Rightarrow \frac{u_x}{u_y} = -\frac{y}{x}.$$

## Maxima & Minima: →

for function of Single Variable: →

$$f(x) \rightarrow \max \rightarrow x = c \text{ if } f(x) \leq f(c) \forall x$$

$$f(x) \rightarrow \min \rightarrow x = c \text{ if } f(x) \geq f(c) \forall x$$

Method: → (i) find  $f'(x)$

(ii) Equate  $f'(x) = 0$  for obtaining the stationary points

(iii) At each stationary pt. find  $f''(x)$

(a) If  $f''(x_0) > 0$  then  $f(x)$  has minima at  $x = x_0$ .

(b) If  $f''(x_0) < 0$  then  $f(x)$  has maxima at  $x = x_0$ .

(c) If  $f''(x_0) = 0$  then  $f(x)$  has no extreme at  $x = x_0$ .

and it is called critical point.

Questions: →

1. The fun  $f(x) = 2x^3 - 3x^2 - 36x + 10$  has a minimum value at  
 $x = \underline{\hspace{2cm}}$ .

- (a) 2      (b)  $\checkmark$  3      (c) -2      (d) -3.

Soln: →  $f'(x) = 6x^2 - 6x - 36 = 0$

$$\Rightarrow x^2 - x - 6 = 0$$

$$\Rightarrow x = -2, 3$$

$$f''(x) = 12x - 6$$

$$f''(x)|_{x=-2} = 12(-2) - 6 = -24 - 6 < 0 \Rightarrow \text{maxima at } x = -2.$$

$$f''(x)|_{x=3} = 12(3) - 6 > 0 \Rightarrow \text{minima at } x = 3.$$

2. The maximum value of fun  $\frac{e^{\sin x}}{e^{\cos x}}$  is \_\_\_\_\_. where,  $x \in R$

(a)  $e^{\frac{1}{\sqrt{2}}}$

(b)  $e^{-\frac{1}{\sqrt{2}}}$

(c)  $e^{\sqrt{2}}$

(d)  $e^{-\frac{1}{\sqrt{2}}}$

25

Sol<sup>n</sup>:--  $f(x) = \frac{e^{\sin x}}{e^{\cos x}} = e^{(\sin x - \cos x)}$

$f(x)$  will have max. value when  $(\sin x - \cos x)$  will be max.

let,  $g(x) = \sin x - \cos x$

$g'(x) = \cos x + \sin x = 0 \Rightarrow x = -\frac{\pi}{4}, \frac{3\pi}{4}$

$g''(x) = -\sin x + \cos x$

$\therefore g''(x) \Big|_{x=-\frac{\pi}{4}} = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} > 0 \Rightarrow$  minima at  $-\frac{\pi}{4}$

$g''(x) \Big|_{x=\frac{3\pi}{4}} = -\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} < 0 \Rightarrow$  maxima at  $\frac{3\pi}{4}$

$\therefore f\left(\frac{3\pi}{4}\right) = e^{\frac{1}{\sqrt{2}} - \left(-\frac{1}{\sqrt{2}}\right)} = e^{\sqrt{2}}$

3. For the fun  $f(x) = x^x$ , minimum appears at  $x =$  \_\_\_\_.

(a)  $e$     (b)  $\frac{1}{e}$     (c)  $e+1$     (d)  $e-1$

Sol<sup>n</sup>:-- let,  $y = x^x \Rightarrow \log y = x \log x$

$\Rightarrow \frac{1}{y} y' = x \cdot \frac{1}{x} + \log x = 1 + \log x$

$\Rightarrow y' = y(1 + \log x) = 0$

$\Rightarrow x = \frac{1}{e}$

4. Consider  $f(x) = (x^2 - 4)^2$  where  $x \in R$ , then,  $f(x)$  has

(a) Only one minima

(b) Only two minima

(c) three minima

(d) three maxima

Soln:  $\rightarrow f(x) = (x^2 - 4)^2$

$$f'(x) = 2 \cdot 2x(x^2 - 4) = 0 \Rightarrow x = \pm 2, 0$$

$$f''(x) = 3x^2 - 4$$

$$f''(0) = -4 < 0 \Rightarrow \text{maxima at } x=0$$

$$f''(2) = 3 \times 4 - 4 > 0 \Rightarrow \text{minima at } x=2$$

$$f''(-2) = 3 \times 4 - 4 > 0 \Rightarrow \text{minima at } x=-2$$

5. If  $f(x) = a \log x + bx^2 - x$  has an extreme value at  $x=-1, 2$

then  $a \& b$  is \_\_\_\_\_.

- (a)  $2, \frac{1}{2}$       (b)  $2, -\frac{1}{2}$       (c)  $-2, \frac{1}{2}$       (d)  $-2, -\frac{1}{2}$

Soln:  $\rightarrow f'(x) = \frac{a}{x} + 2bx - 1 = 0$

$$\Rightarrow a + 2bx^2 - x = 0$$

$$\Rightarrow 2bx^2 - x + a = 0$$

$$\Rightarrow x^2 - \frac{1}{2b}x + \frac{a}{2b} = 0$$

$$\therefore \frac{1}{2b} = -1 + 2 = 1 \Rightarrow b = \frac{1}{2}$$

$$\frac{a}{2b} = -2 \Rightarrow a = -2$$

6. The maximum value of  $f(x) = x^2 - x - 2$  in  $[-4, 4]$  is \_\_\_\_\_.

- (a) 18      (b) 10      (c) -2.25      (d) indeterminate

Soln:  $\rightarrow f'(x) = 2x - 1 = 0 \Rightarrow x = \frac{1}{2}$

$$f''(x) = 2 > 0 \Rightarrow \text{minima at } x = \frac{1}{2}$$

$$f(-4) = 18$$

$$f(4) = 10$$

## Maxima & Minima for fun. of two variables:

26

let  $z = f(x, y)$ ,

consider,  $P = \frac{\partial z}{\partial x}$ ,  $Q = \frac{\partial z}{\partial y}$ ,  $R = \frac{\partial^2 z}{\partial x^2}$ ,  $S = \frac{\partial^2 z}{\partial x \partial y}$ ,  $T = \frac{\partial^2 z}{\partial y^2}$

Method: → (i) Find  $P, Q, R, S$  &  $T$

(ii) Equate  $P$  &  $Q$  to zero for obtaining the stationary points

(iii) At each stationary point find  $R, S$  &  $T$ .

- (a) If  $RT - S^2 > 0$ ,  $R > 0$  then the fun  $f(x, y)$  has a minima at that stationary point.
- (b) If  $RT - S^2 > 0$ ,  $R < 0$  then  $f(x, y)$  has a maxima at that stationary point.
- (c) If  $RT - S^2 < 0$ , then  $f(x, y)$  has no extreme at that stationary point & it is known as Saddle point

### Questions:

1. The fun  $f(x, y) = x^2 + y^2 + 6x = 0$  has

- (a) min. at  $(-3, 0)$
- (b) max. at  $(-3, 0)$
- (c)  $(-3, 0)$  is a saddle point
- (d) none

### Sol<sup>n</sup>:

2. The fun  $f(x,y) = x^3 - 3x^2 + 4y^2 + 6$  has a minimum value at  $x = \underline{\hspace{2cm}}$ .

- (a) (0, 0)      (b) (2, 0)      (c) (2, 1)      (d) (-2, 0)

$$\text{Soln: } \rightarrow P = \frac{\partial f}{\partial x} = 3x^2 - 6x = 0 \Rightarrow x = 0, 2.$$

$$Q = \frac{\partial f}{\partial y} = 8y = 0 \Rightarrow y = 0$$

$$R = \frac{\partial^2 f}{\partial x^2} = 6x - 6$$

$$S = \frac{\partial^2 f}{\partial x \partial y} = 0$$

$$T = \frac{\partial^2 f}{\partial y^2} = 8$$

At (0, 0) & (2, 0)

$$R = -6 < 0$$

$$S = 0$$

$$T = 8$$

$$RT - S^2 < 0$$

Saddle point

$$R = 6 > 0$$

$$S = 0$$

$$T = 8$$

$$RT - S^2 > 0$$

minima at (2, 0)

3. The fun  $f(x,y) = x^3 + y^3 - 3axy$  has

(a) max. at (a, a)

(b) max. at (a, a) if  $a < 0$

(c) min. at (a, a)

(d) max. at (a, a) if  $a > 0$

$$\text{Soln: } \rightarrow P = 3x^2 - 3ay = 0 \Rightarrow x^2 = ay \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{solving}$$

$$Q = 3y^2 - 3ax = 0 \Rightarrow y^2 = ax$$

Q. 4: A rectangular box open at the top is to have a volume of  $32 \text{ ft}^3$ , then, the dimensions of the box requiring the least material for its construction is \_\_\_\_.

- (a) 8, 2, 2      (b)  $\checkmark$  4, 4, 2      (c) 16, 1, 2      (d) 8, 8,  $\frac{1}{2}$

Soln: Let the dimension of the box is  $x, y, z \Rightarrow xyz = 32$

$$\therefore S = xy + 2yz + 2xz$$

$$\text{i.e. } f(x, y) = xy + 2y \cdot \frac{32}{xy} + 2x \cdot \frac{32}{xy}$$

$$= xy + \frac{64}{x} + \frac{64}{y}$$

$$P = y - \frac{64}{x^2} = 0 \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{Solving } x=4, y=4$$

$$q = x - \frac{64}{y^2} = 0$$

Q. 5. The distance between the origin and a point nearest to it on the surface  $z^2 = 1+xy$  is \_\_\_\_.

- (a)  $\sqrt{3}$       (b)  $\sqrt{2}$       (c)  $\checkmark$  1      (d) None

Soln: Let  $P(x, y, z)$  be on the surface  $z^2 = 1+xy$

$$\therefore D = \sqrt{x^2 + y^2 + z^2} = \sqrt{x^2 + y^2 + xy + 1}$$

Consider,  $f = x^2 + y^2 + xy + 1$

$$P = 2x + y = 0 \quad \left. \begin{array}{l} \\ \end{array} \right\} \text{Solving } x=0, y=0$$

$$Q = 2y + x = 0$$

$$x=2$$

$$x=1$$

$$t=2$$

$$y=t-s^2 = 4-1 = 3 > 0 \text{ also } t>0$$

i.e minima at  $x=0, y=0$

at  $x=0, y=0$  we have  $z^2 = 1+0 = 1 \Rightarrow z = \pm 1$

$$\therefore D = \sqrt{1} = 1.$$

## Constrained Maxima & Minima:

### Lagrange's method of undetermined multipliers:

Let  $f(x, y, z)$  &  $\phi(x, y, z) = c$ , then

$$F(x, y, z) = f(x, y, z) + \lambda \phi(x, y, z) = 0$$

$$F_x = 0 \quad \text{i.e.} \quad \frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0$$

$$F_y = 0 \quad \text{i.e.} \quad \frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0$$

$$F_z = 0 \quad \text{i.e.} \quad \frac{\partial f}{\partial z} + \lambda \frac{\partial \phi}{\partial z} = 0$$

} Lagrange's eqn

$\lambda \rightarrow$  Lagrange's Multiplier

### Questions:

1. The volume of the greatest parallelopiped that can be inscribed in an

ellipse:  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$  is \_\_\_\_\_.

- (a)  $\frac{8abc}{3\sqrt{3}}$     (b)  $\frac{4abc}{3\sqrt{3}}$     (c)  $\frac{abc}{\sqrt{3}}$     (d)  $\frac{abc}{3\sqrt{3}}$

Soln: Let  $2x, 2y, 2z$  be dimension of parallelopiped,

$$\text{volume} = 8xyz = f(x, y, z)$$

$$\phi(x, y, z) = \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 = 0$$

$$\therefore F(x, y, z) = 8xyz + \lambda \left( \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 \right) = 0$$

$$F_x = 8yz + \frac{2\lambda x}{a^2} = 0 \Rightarrow -\frac{\lambda}{4} = \frac{a^2yz}{x}$$

$$F_y = 8xz + \frac{2\lambda y}{b^2} = 0 \Rightarrow -\frac{\lambda}{4} = \frac{b^2xz}{y}$$

$$F_z = 8xy + \frac{2\lambda z}{c^2} = 0 \Rightarrow -\frac{\lambda}{4} = \frac{c^2xy}{z}$$

$$\therefore \frac{a^2yz}{x} = \frac{b^2xz}{y} \quad \& \quad \frac{b^2xz}{y} = \frac{c^2xy}{z} \quad \& \quad \frac{c^2xy}{z} = \frac{a^2yz}{x}$$

$$\Rightarrow \frac{x^2}{a^2} = \frac{y^2}{b^2} \quad \& \quad \frac{y^2}{b^2} = \frac{z^2}{c^2} \quad \& \quad \frac{z^2}{c^2} = \frac{x^2}{a^2}$$

$$\Rightarrow \frac{3x^2}{a^2} = 1 \Rightarrow x = \frac{a}{\sqrt{3}} \quad \& \quad y = \frac{b}{\sqrt{3}}, z = \frac{c}{\sqrt{3}}$$

$$\therefore \text{volume, } V = 8xyz = \frac{8abc}{3\sqrt{3}}.$$

28

2. The min value of  $x^2+y^2+z^2$  where  $x+y+z=1$  is \_\_\_\_.

- (A)  $\frac{1}{3}$     (B)  $\frac{1}{9}$     (C)  $\frac{1}{27}$     (D) 1

Soln:  $\rightarrow f = (x^2+y^2+z^2) \text{ & } \phi = x+y+z-1$

$$\therefore F = f + \lambda \phi$$

$$\Rightarrow F_x = 2x + \lambda \Rightarrow -\lambda = 2x$$

$$F_y = 2y + \lambda \Rightarrow -\lambda = 2y$$

$$F_z = 2z + \lambda \Rightarrow -\lambda = 2z$$

$$\Rightarrow x=y=z$$

$$\therefore x = \frac{1}{3}, y = \frac{1}{3}, z = \frac{1}{3}$$

$$\therefore f_{\min} = x^2+y^2+z^2 = 3x^2 = \frac{1}{3}$$

### 5. Multiple Integrals: $\rightarrow$


Double Integral:  $\rightarrow$  Let  $f(x,y)$  be defined at each point in the given region  $R$ , then its double integral is,  $\iint_R f(x,y) dx dy$ , where, continuity of  $f(x,y)$  in the region  $R$  ensures the existence of the integral.

#### Methods of Evaluation: $\rightarrow$

Case 1:  $\rightarrow$  Let the limits of integration be  $y=\phi_1(x)$  to  $y=\phi_2(x)$  &

$x=c_1$  to  $c_2$ .

$$\iint_R f(x,y) dx dy = \int_{c_1}^{c_2} \left[ \int_{\phi_1(x)}^{\phi_2(x)} f(x,y) dy \right] dx$$




$c_1$  to  $c_2$  to get area

Case 2— when the limits of integration are  $x = \phi_1(y)$  &  $x = \phi_2(y)$ , and

$y = d_1$  to  $y = d_2$ ,


$$\iint_R f(x,y) dx dy = \int_{y=d_1}^{d_2} \left[ \int_{x=\phi_1(y)}^{\phi_2(y)} f(x,y) dx \right] dy$$



Case 3— when the limits are  $x = c_1$  to  $x = c_2$  &  $y = d_1$  to  $y = d_2$ ,

$$\iint_R f(x,y) dx dy = \int_{x=c_1}^{c_2} \left[ \int_{y=d_1}^{d_2} f(x,y) dy \right] dx$$

$$= \int_{y=d_1}^{d_2} \left[ \int_{x=c_1}^{c_2} f(x,y) dx \right] dy$$



Questions:→

1. Evaluate the following

$$(i) \int_0^1 \int_0^2 (xy + x^3) dx dy. \quad (ii) \int_0^4 \left[ \int_0^{x^2} e^{yx} dy \right] dx.$$

$$\text{Soln: } (i) \int_0^1 \left( \frac{x^2}{2}y + \frac{x^4}{4} \right)_0^2 dy = \int_0^1 (2y + 4) dy = [y^2 + 4y]_0^1 = 5.$$

$$(ii) \int_0^4 \left[ \frac{e^{yx}}{1/x} \right]_0^{x^2} dx = \int_0^4 (x e^{x^2} - x) dx = \left[ x e^{x^2} - e^x - \frac{x^2}{2} \right]_0^4 \\ = [4e^4 - e^4 - 8] - [-1] = 3e^4 - 7.$$

2. The value of  $\iint_R xy dx dy$ , where, R is a region in the 1st +ve quad-

-rant at the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is \_\_\_\_\_.

- (a)  $\frac{a^2 b^2}{8}$     (b)  $\frac{ab}{8}$     (c)  $\frac{a^3 b^3}{8}$     (d)  $\frac{a^2 b^2}{4}$

24

Sol<sup>n</sup>: Consider the horizontal strip,

$$x=0 \text{ to } x=\frac{a}{b} \sqrt{b^2-y^2}$$


$$y=0 \text{ to } y=b$$

$$\therefore \iint_R xy \, dx \, dy = \int_{y=0}^b \int_{x=0}^{\frac{a}{b} \sqrt{b^2-y^2}} xy \, dx \, dy$$

$$= \int_{y=0}^b \left[ \frac{x^2 y}{2} \right]_0^{\frac{a}{b} \sqrt{b^2-y^2}} \, dy$$

$$= \int_0^b \frac{\frac{a^2}{b^2} (b^2-y^2) y}{2} \, dy$$

$$= \left[ \frac{a^2}{2} \cdot \frac{y^2}{2} - \frac{a^2}{2b^2} \cdot \frac{y^4}{4} \right]_0^b = \frac{a^2 b^2}{4} + \frac{a^2 b^2}{8} = \frac{a^2 b^2}{8}$$



3. The value of  $\iint_R y \, dx \, dy$ , where R is a region  $y=x^2$ ,  $x+y=2$

$x=0$  is \_\_\_\_\_.

$$\underline{\text{Sol<sup>n</sup>:}} \quad x+x^2=2$$

$$\Rightarrow x^2+x-2=0$$

$$\Rightarrow (x-1)(x+2)=0$$

$$\Rightarrow x=1, -2$$


$\therefore$  pt. of intersection is  $(1,1)$

Consider the vertical strip,

$$y=x^2 \text{ to } y=2-x$$

$$x=0 \text{ to } 1$$

$$\therefore \iint_R y \, dx \, dy = \int_{x=0}^1 \int_{y=x^2}^{2-x} y \, dy \, dx = \int_0^1 \left[ \frac{y^2}{2} \right]_{x^2}^{2-x} \, dx = \frac{16}{15}$$



Ques-4. The value of  $\iint_R r^2 \sin\theta dr d\theta$ , where,  $R$  is the region, bounded by the semicircle  $r = 2a \cos\theta$  above the initial line is \_\_\_\_\_.

Sol<sup>n</sup>:  $\rightarrow r = 0$  to  $2a \cos\theta$

$\theta = 0$  to  $\pi/2$


$$\iint_R r^2 \sin\theta dr d\theta$$

$$= \int_{\theta=0}^{\pi/2} \int_{r=0}^{2a \cos\theta} r^2 \sin\theta dr d\theta$$

$$= \int_0^{\pi/2} \sin\theta \left[ \frac{r^3}{3} \right]_0^{2a \cos\theta} d\theta$$

$$= \int_0^{\pi/2} \frac{8a^3 \cos^3\theta \sin\theta}{3} d\theta$$

$$= -\frac{8a^3}{3} \left[ \frac{\cos^4\theta}{4} \right]_0^{\pi/2} = \frac{2a^3}{3}$$



Change of order of integration:  $\rightarrow$

Questions:  $\rightarrow$

1. The value of  $\int_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx$  is \_\_\_\_\_.


Sol<sup>n</sup>: Given limits are  $y=x$  to  $y=\infty$   
 $x=0$  to  $x=\infty$

Horizontal Strip: —

$$x=0, x=y$$

$$y=0, y=\infty$$

$$\iint_0^\infty \int_x^\infty \frac{e^{-y}}{y} dy dx = \int_{y=0}^\infty \left[ \int_{x=0}^y \frac{e^{-y}}{y} dx \right] dy = \int_0^\infty \left[ \frac{e^{-y} y}{y} \right]_0^\infty dy = \int_0^\infty e^{-y} dy$$



Q. 2. By reversing the order of integration  $\int_0^2 \int_{x^2}^{2x} f(x,y) dy dx$  may be represented as.

$$(a) \int_0^2 \int_{x^2}^{2x} f(x,y) dy dx$$

$$(b) \int_0^2 \int_y^{2x} f(x,y) dx dy$$

$$(c) \int_0^4 \int_{y/2}^{\sqrt{y}} f(x,y) dx dy$$

$$(d) \int_{x^2}^{2x} \int_0^2 f(x,y) dx dy$$

Soln: Given limits are


$$x=0 \text{ to } x=2$$

$$y=x^2 \text{ to } y=2x$$

Horizontal Strip:

$$y=0 \text{ to } y=4$$

$$x=\frac{y}{2} \text{ to } x=\sqrt{y}$$



3. By changing the order of integration  $\int_0^8 \int_{x/4}^2 f(x,y) dy dx$  leads to

$$\int_p^q \int_0^{x/4} f(x,y) dx dy \text{ then } q \text{ is } \underline{\hspace{2cm}}$$

Soln: Given limits are

$$y=\frac{x}{4} \text{ to } y=2$$


$$x=0 \text{ to } x=8$$

By changing limits,

$$y=0 \text{ to } y=2$$

$$x=0 \text{ to } x=4y$$

$$q=4y$$



Triple Integrals → Let  $f(x, y, z)$  be defined at each point in the region 'R' of space then its triple integral is  $\iiint_R f(x, y, z) dx dy dz$ .

Let  $z = \phi_1(x, y)$  to  $z = \phi_2(x, y)$

$y = \psi_1(x)$  to  $y = \psi_2(x)$

$x = c_1$  to  $x = c_2$ , then

$$\iiint_R f(x, y, z) dx dy dz = \int_{x=c_1}^{c_2} \int_{y=\psi_1(x)}^{\psi_2(x)} \int_{z=\phi_1(x, y)}^{\phi_2(x, y)} f(x, y, z) dz dy dx$$

Questions →

1. Evaluate  $\int_0^2 \int_0^x \int_0^{\sqrt{x+y}} z dz dy dx$ .

Soln: →  $I = \int_0^2 \int_0^x \left[ \frac{z^2}{2} \right]_0^{\sqrt{x+y}} dy dx = \int_0^2 \left[ \frac{(x+y)^2}{4} \right]_0^x dx = 2$ .

2. The value of  $\iiint_R y dx dy dz$ , where R is the region bounded by

the plane  $x=0, y=0, z=0$  &  $x+y+z=1$  is

Soln: →  $\int_0^1 \int_0^{1-x} \int_0^{1-x-y} y dz dy dx = \int_0^1 \int_0^{1-x} y(1-x-y) dy dx$

$$= \int_0^1 \left[ y - xy - \frac{y^2}{2} \right]_0^{1-x} dx$$

$$= \int_0^1 \left[ 1-x - x+x^2 - \frac{(1-x)^2}{2} \right] dx$$

$$= \int_0^1 \left[ \frac{(1-x)y^2}{2} - \frac{y^3}{3} \right]_0^{1-x} dx$$

$$= \int_0^1 \left[ \frac{(1-x)^3}{2} - \frac{(1-x)^3}{3} \right] dx = \frac{(1-x)^4}{24} \Big|_0^1 = \frac{1}{24}$$

Change of variables :→ In a double integral, if  $x = f(u, v)$  &  $y = g(u, v)$

then,  $\iint_R \phi(x, y) dx dy = \iint_R \phi(f, g) |J| du dv = \iint_R \phi(u, v) |J| du dv$

where,  $|J| \rightarrow$  Jacobian of transformation used to transform one system to another.

$$|J| = J\left(\frac{x, y}{u, v}\right) = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

Cartesian form → Polar form :→

$$(x, y) \rightarrow (r, \theta)$$

$$x = r \cos \theta, \quad y = r \sin \theta$$

$$|J| = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r$$

$$\Rightarrow \iint_R \phi(x, y) dx dy = \iint_R \psi(r, \theta) r dr d\theta$$

In a triple integral, if  $x = f(u, v, w)$ ,  $y = g(u, v, w)$  &  $z = h(u, v, w)$

then,

$$\iiint_R f(x, y, z) dx dy dz = \iiint_R \psi(u, v, w) |J| du dv dw$$

where,  $|J| = J\left(\frac{x, y, z}{u, v, w}\right) = \frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$

Cartesian to Cylindrical Polar Form : →

$$(x, y, z) \quad (r, \theta, z)$$

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z$$

$$|J| = \begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$$

$$\iiint_R \phi(x, y, z) dx dy dz = \iiint_R \psi(r, \theta, z) r dr d\theta dz$$

Cartesian to Spherical Polar Form : →

$$(x, y, z) \quad (r, \theta, \phi)$$

$$x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta$$

$$|J| = \begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix} = r^2 \sin \theta$$

$$\iiint_R \phi(x, y, z) dx dy dz = \iiint_R \psi(r, \theta, \phi) r^2 \sin \theta dr d\theta d\phi$$

Questions : →


1. Evaluate  $\int_0^\infty \int_0^\infty e^{-(x^2+y^2)} dx dy$

Soln : → let  $x = r \cos \theta, \quad y = r \sin \theta$ , then  $|J| = r$

$$\therefore x^2 + y^2 = r^2$$

$$\therefore \int_0^\infty \int_0^{\pi/2} e^{-r^2} \cdot r dr d\theta$$

let  $r^2 = t \Rightarrow r dr = \frac{dt}{2}$



$$0 \rightarrow 0 \text{ to } \frac{\pi}{2}$$

$$\Rightarrow \int_0^{\pi/2} \int_0^\infty e^{-t} \frac{dt}{2} d\theta = \int_0^{\pi/2} \left[ \frac{e^{-t}}{2} \right]_0^\infty d\theta = \frac{1}{2} \int_0^{\pi/2} d\theta = \frac{\pi}{4}. \quad 82$$

2. Evaluate  $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} \frac{dz dy dx}{\sqrt{1-x^2-y^2-z^2}}$ .

Sol<sup>n</sup>:  $\rightarrow z=0 \text{ to } z=\sqrt{1-x^2-y^2} \Rightarrow z^2=1-x^2-y^2 \Rightarrow x^2+y^2+z^2=1$

Using spherical coordinates,

$$x = r \sin\theta \cos\phi, \quad y = r \sin\theta \sin\phi, \quad z = r \cos\theta$$

$$x^2+y^2+z^2 = r^2$$

$$\Rightarrow r \rightarrow 0 \text{ to } 1:$$

$$\theta \rightarrow 0 \text{ to } \pi/2$$

$$\phi \rightarrow 0 \text{ to } \pi/2$$

$$\begin{aligned} \int_0^{\pi/2} \int_0^{\pi/2} \int_0^1 \frac{r^2 \sin\theta dr d\theta d\phi}{\sqrt{1-r^2}} &= \int_0^{\pi/2} \int_0^{\pi/2} \int_0^1 \sin\theta \left[ \frac{1}{\sqrt{1-r^2}} - \sqrt{1-r^2} \right] dr d\theta d\phi \\ &= \int_0^{\pi/2} \int_0^{\pi/2} \sin\theta \left[ \sin^{-1} r - \left( \frac{r \sqrt{1-r^2}}{2} + \frac{1}{2} \sin^{-1}(r) \right) \right]_0^1 d\theta d\phi \\ &= \int_0^{\pi/2} \int_0^{\pi/2} \sin\theta \left[ \frac{\pi}{4} \right] d\theta d\phi \\ &= \frac{\pi}{4} \int_0^{\pi/2} [-\cos\theta]_0^{\pi/2} d\phi = \frac{\pi}{4} \cdot \frac{\pi}{2} = \frac{\pi^2}{8}. \end{aligned}$$

3. By a change of variable  $x(u,v) = uv$  &  $y(u,v) = \frac{v}{u}$  in double integral  
the integrand  $f(x,y)$  changes to  $f(uv, \frac{v}{u}) \phi(u,v)$  then  $\phi(u,v)$  is

- (a)  $\frac{2v}{u}$    (b)  $2uv$    (c)  $v^2$    (d)  $= 1$

Sol<sup>n</sup>:  $\phi(u,v) = |J| = \begin{vmatrix} v & u \\ -\frac{v}{u^2} & \frac{1}{u} \end{vmatrix} = \frac{v}{u} + \frac{v}{u} = \frac{2v}{u}.$

### Lengths, Areas & Volumes:

(i) Length of an arc of a curve  $y=f(x)$  between the lines  $x=x_1$  &  $x=x_2$

$$\text{is, } l = \int_{x_1}^{x_2} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

(ii) length of an arc of a curve  $x=f(t)$  &  $y=g(t)$  between  $t=t_1$  to  $t=t_2$

$$\text{is, } l = \int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

(iii) Area of the region bounded by the curve  $y=f(x)$  &  $y=g(x)$  between

$x=x_1$  &  $x=x_2$  is,

$$A = \int_{x_1}^{x_2} [g(x) - f(x)] dx \quad (\text{or}) \quad \int_{x_1}^{x_2} \int_{f(x)}^{g(x)} dy dx$$

(iv) The volume of solid generated by revolving  $y=f(x)$  between  $x=x_1$  &  $x=x_2$  about  $x$ -axis is,

$$V = \int_{x_1}^{x_2} \pi y^2 dx$$

$$\text{about } y\text{-axis is, } V = \int_{y_1}^{y_2} \pi x^2 dy$$

In polar form:- (i) about initial line (i.e.  $\theta=0^\circ$ )

$$V = \int_{\theta_1}^{\theta_2} \frac{2\pi}{3} r^3 \sin\theta d\theta$$

(ii) about line  $\theta=\frac{\pi}{2}$

$$V = \int_{\theta_1}^{\theta_2} \frac{2\pi}{3} r^3 \cos\theta d\theta$$

Questions:

33

1. The length of the curve  $y = \frac{2}{3}x^{3/2}$  between  $x=0$  &  $1$  is — .  
 (a) 0.27    (b) 0.67    (c) 1    (d) 1.22

Sol<sup>n</sup>:—  $\frac{dy}{dx} = \frac{2}{3} \cdot \frac{3}{2} x^{\frac{3}{2}-1} = x^{1/2}$

$$\therefore l = \int_{x_1}^{x_2} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^1 \sqrt{1+x} dx = \left[ \frac{(1+x)^{3/2}}{3/2} \right]_0^1 = 1.22$$

2. The length of the curve,  $x = \cos^3\theta$ ,  $y = \sin^3\theta$  between  $\theta=0$  &  $\pi/2$  is —

Sol<sup>n</sup>:— 
$$l = \int_0^{\pi/2} \sqrt{(3\cos^2\theta(-\sin\theta))^2 + (3\sin^2\theta\cos\theta)^2} d\theta$$

$$= \int_0^{\pi/2} (3\sin\theta\cos\theta)\sqrt{\sin^2\theta + \cos^2\theta} d\theta = \frac{3}{2}$$

3. The area bounded by the parabola  $y = 2x^2$  & st. line  $x = y - 4$ .

Ans: —


Sol<sup>n</sup>:— Pt. of intersection—

$$y = 2(y-4)^2$$

$$\Rightarrow y = 2(y^2 + 16 - 8y)$$

$$\Rightarrow 2y^2 - 17y + 32 = 0$$

$$\Rightarrow y = \frac{17 \pm \sqrt{289-256}}{4} = \frac{17 \pm \sqrt{33}}{4}$$




$$A = \int_{\frac{17-\sqrt{33}}{4}}^{\frac{17+\sqrt{33}}{4}} [(y-4) - 2x^2] dy$$

$$A = \int_{\frac{17-\sqrt{33}}{4}}^{\frac{17+\sqrt{33}}{4}} [(y-4) - \frac{y^2}{2}] dy$$

4. The area between the curves  $y^2 = 4x$  &  $x^2 = 4y$  is, —

Sol<sup>n</sup>:—  $A = \int_0^4 (2\sqrt{x} - \frac{x^2}{4}) dx = \frac{16}{3}$ .



5. The volume generated by revolving the area bounded by parabola  $y^2 = 8x$  and st. line  $x=2$  about y-axis is \_\_\_\_\_.

(a)  $\frac{128\pi}{5}$

(b)  $\frac{5\pi}{128}$

(c)  $\frac{127\pi}{5}$

(d) none


Sol<sup>n</sup>:  $V = \int_{y_1}^{y_2} \pi x^2 dy$

$$= \int_{-4}^4 \pi \frac{y^4}{64} dy$$

$$= \frac{\pi}{64} \left[ \frac{y^5}{5} \right]_{-4}^4 = \frac{32\pi}{5}$$

$$\text{Total volume} = \int_{-4}^4 \pi x^2 dy = \int_{-4}^4 \pi (2)^2 dy = 4\pi [y]_{-4}^4 = 32\pi$$

$$\therefore \text{Remaining volume} = 32\pi - \frac{32\pi}{5} = \frac{128\pi}{5}$$



6. The volume of solid generated by revolving the cardiac  $r = a(1-\cos\theta)$  about the initial line is \_\_\_\_\_.


Sol<sup>n</sup>:  $V = \int_{\theta_1}^{\theta_2} \frac{2\pi}{3} r^3 \sin\theta d\theta$

$$= \int_0^{\pi} \frac{2\pi}{3} a^3 (1-\cos\theta)^3 \sin\theta d\theta$$

$$\text{Set } 1-\cos\theta = t \Rightarrow \sin\theta d\theta = dt$$

$$\text{as } \theta = 0, t = 0$$

$$\theta = \pi, t = 2$$



$$\therefore V = \int_0^2 \frac{2\pi}{3} a^3 t^3 dt = \frac{2\pi a^3}{3} \left[ \frac{t^4}{4} \right]_0^2 = \frac{8\pi a^3}{3}$$

### Vector Calculus:

Position Vector: → The position vector of the point  $P(x, y, z)$  in the space is

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}, \quad |\vec{r}| = \sqrt{x^2 + y^2 + z^2}$$

In parametric form,

$$\vec{r} = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$$

$$\text{let } \vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

$$(i) \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos(\vec{a}, \vec{b}) = a_1 b_1 + a_2 b_2 + a_3 b_3$$

(ii)  $\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin(\vec{a}, \vec{b}) \hat{n}$  where  $\hat{n}$  is vector of unit length perpendicular


to the plane contains  $\vec{a}$  &  $\vec{b}$ .

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Note: → (i) Area of  $\triangle OAB$

$$= \frac{1}{2} |\vec{OA} \times \vec{OB}|$$


$$= \frac{1}{2} |\vec{a} \times \vec{b}|$$



(ii) Area of  $\triangle ABC$


$$= \frac{1}{2} |\vec{AB} \times \vec{AC}|$$

$$= \frac{1}{2} |(\vec{b} - \vec{a}) \times (\vec{c} - \vec{a})|$$



(iii) Area of parallelogram

$$= |\vec{a} \times \vec{b}|$$



### Scalar Triple Products:

$$(\vec{a} \times \vec{b}) \cdot \vec{c} = \vec{a} \cdot (\vec{b} \times \vec{c}) = [\vec{a} \vec{b} \vec{c}]$$

$$= \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

Note: → (i) Volume of parallelopiped,  $V = |[\vec{a} \vec{b} \vec{c}]|$

(ii) Volume of tetrahedron,  $V = \frac{1}{6} |[\vec{a} \vec{b} \vec{c}]|$

where,  $\vec{a}$ ,  $\vec{b}$  &  $\vec{c}$  are edge vectors of parallelopiped parallelopip.

Vector Triple Product: →

$$(\vec{a} \times \vec{b}) \times \vec{c} = (\vec{c} \cdot \vec{a}) \vec{b} - (\vec{c} \cdot \vec{b}) \vec{a}$$

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{c} \cdot \vec{a}) \vec{b} - (\vec{b} \cdot \vec{a}) \vec{c}$$

6. Vector Differentiation: → Let  $\vec{x}(t) = \vec{f}(t)$  be the vector eqn of the curve

then  $\frac{d\vec{r}}{dt} = \lim_{\Delta t \rightarrow 0} \frac{\vec{f}(t + \Delta t) - \vec{f}(t)}{\Delta t}$

If  $t$  is a time variable then  $\frac{d\vec{r}}{dt}$  represents a velocity vector.

Note: → (i)  $\frac{d\vec{r}}{dt}$  is a vector in the direction of tangent to the curve at that point.

(ii)  $\vec{F}(t)$  is constant in magnitude then,  $\vec{F} \cdot \frac{d\vec{F}}{dt} = 0$

(iii)  $\vec{F}(t)$  vector with fixed direction then,  $\vec{F} \times \frac{d\vec{F}}{dt} = 0$

Properties: → Let  $\vec{a}(t)$  &  $\vec{b}(t)$  be the vector fun of the scalar variable 't' and  $\phi$  be a scalar fun then,

(i)  $\frac{d}{dt} (\vec{a} \pm \vec{b}) = \frac{d\vec{a}}{dt} \pm \frac{d\vec{b}}{dt}$

(ii)  $\frac{d}{dt} (\vec{a} \cdot \vec{b}) = \frac{d\vec{a}}{dt} \cdot \vec{b} + \vec{a} \cdot \frac{d\vec{b}}{dt}$

(iii)  $\frac{d}{dt} (\vec{a} \times \vec{b}) = \left( \frac{d\vec{a}}{dt} \times \vec{b} \right) + \left( \vec{a} \times \frac{d\vec{b}}{dt} \right)$

(iv)  $\frac{d}{dt} (\phi \vec{a}) = \frac{d\phi}{dt} \vec{a} + \phi \frac{d\vec{a}}{dt}$

Point function: $\rightarrow$  If the value of function depends on position of point, then it is said to be point fun.

35

Scalar Point function: $\rightarrow$  If to each point  $P(x,y,z)$  in region  $R$  of space  $\exists$  a unit scalar associated with it, then,  $\phi(x,y,z)$  is a scalar point function.

The set of all points in the region  $R$  of space together with the scalar values forms a scalar field.

$\&$ : $\rightarrow$  The temp.  $T(x,y,z)$  at any point on a body is a scalar point function and the medium it self is a scalar field.

Vector Point Function: $\rightarrow$  The velocity at any time  $t$  of a particle in a fluid flow is a vector point fun.

Vector Differential Operator: $\rightarrow$  (del)  $\vec{\nabla}$

$$\vec{\nabla} = \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z}$$

Level Surface: $\rightarrow$  Let  $\phi(x,y,z)$  be a scalar field in the region ' $R$ ', then the set of points satisfying  $\phi(x,y,z) = c$ , where, ' $c$ ' is an arbitrary const, constitutes a family of surfaces called level surfaces.

Gradient of a Scalar function: $\rightarrow$  Let  $\phi(x,y,z)$  be a differentiable scalar pt. fun then gradient of scalar denoted by grad  $\phi$

$$(or) \vec{\nabla} \phi = \hat{i} \frac{\partial \phi}{\partial x} + \hat{j} \frac{\partial \phi}{\partial y} + \hat{k} \frac{\partial \phi}{\partial z}$$

$\hookrightarrow$  vector normal to surface  $\phi$

$$\frac{\vec{\nabla} \phi}{|\vec{\nabla} \phi|} \rightarrow$$
 unit vector normal to surface  $\phi$ .

$$(i) \vec{\nabla} \cdot \vec{F} = \vec{F} \cdot \vec{\nabla}$$

Questions: →

1.  $\vec{\nabla} r$  is \_\_\_\_\_. if  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

Sol<sup>n</sup>:→  $r = \sqrt{x^2 + y^2 + z^2} \Rightarrow \vec{\nabla} r = \frac{\vec{r}}{|\vec{r}|}$

Note:→  $\vec{\nabla} f(r) = f'(r) \frac{\vec{r}}{r}$

Eg:→  $\vec{\nabla} (\log r) = \frac{1}{r} \cdot \frac{\vec{r}}{r} = \frac{\vec{r}}{r^2}$

$\vec{\nabla} (\sin \log r) = \frac{\cos \log r}{r} \cdot \frac{\vec{r}}{r} = \cos \log r \cdot \frac{\vec{r}}{r^2}$

2. The unit vector normal to the surface  $y^2 = 8x$  at  $(1, 2)$  is \_\_\_\_.

Sol<sup>n</sup>:→ Let  $\phi = y^2 - 8x$

$\vec{\nabla} \phi = -8\hat{i} + 2y\hat{j}$

$\vec{\nabla} \phi|_{(1,2)} = -8\hat{i} + 4\hat{j}$

$\therefore \hat{n} = \frac{\vec{\nabla} \phi}{|\vec{\nabla} \phi|} = \frac{-8\hat{i} + 4\hat{j}}{\sqrt{64+16}} = \frac{-8\hat{i} + 4\hat{j}}{\sqrt{80}}$

unit

3. A sphere of radius is centred at origin. The unit normal at a point  $P(x, y, z)$  to the surface of the sphere is the vector

- (a)  $(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$    (b)  $(\frac{x}{\sqrt{3}}, \frac{y}{\sqrt{3}}, \frac{z}{\sqrt{3}})$    (c)  $(x, y, z)$    (d)  $(\frac{x}{\sqrt{2}}, \frac{y}{\sqrt{2}}, \frac{z}{\sqrt{2}})$

Sol<sup>n</sup>:→  $x^2 + y^2 + z^2 = 1$

Let  $\phi = x^2 + y^2 + z^2$

$\vec{\nabla} \phi = 2x\hat{i} + 2y\hat{j} + 2z\hat{k}$

$\frac{\vec{\nabla} \phi}{|\vec{\nabla} \phi|} = \frac{2x\hat{i} + 2y\hat{j} + 2z\hat{k}}{\sqrt{4x^2 + 4y^2 + 4z^2}} = x\hat{i} + y\hat{j} + z\hat{k}$

Directional Derivative: $\rightarrow$  The directional derivative of a differentiable scalar fun  $\phi(x, y, z)$  in the direction of  $\vec{a}$  is given by,

$$D.D. = \vec{\nabla} \phi \cdot \frac{\vec{a}}{|\vec{a}|}$$

Let,  $\vec{a} = \hat{i}$ , then,

$$\begin{aligned} D.D. &= \vec{\nabla} \phi \cdot \frac{\hat{i}}{|\hat{i}|} = \left( \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k} \right) \cdot \hat{i} \\ &= \frac{\partial \phi}{\partial x}. \end{aligned}$$

Let,  $\frac{\vec{a}}{|\vec{a}|} = \hat{b}$ , then,  $D.D. = \vec{\nabla} \phi \cdot \hat{b} = |\vec{\nabla} \phi| |\hat{b}| \cos \theta = |\vec{\nabla} \phi| \cos \theta$

The max. value of  $\cos \theta = 1$  i.e. when  $\theta = 0$  i.e. when  $\hat{b}$  is parallel to  $\vec{\nabla} \phi$ . Therefore, the value of the directional derivative is maximum in the direction of normal to the surface  $\phi$ , and the maximum value of directional derivative is  $|\vec{\nabla} \phi|$ .

Angle Between Surfaces: $\rightarrow$  Angle between the normal to the surfaces at the pt. of intersection is taken as the angle between the surfaces. Let  $\theta$  be the angle b/w the surfaces  $\phi_1(x, y, z) = c_1$  &  $\phi_2(x, y, z) = c_2$  then

$$\cos \theta = \frac{\vec{\nabla} \phi_1 \cdot \vec{\nabla} \phi_2}{|\vec{\nabla} \phi_1| |\vec{\nabla} \phi_2|}$$

Questions:

1. The directional derivative of  $f(x, y) = x^2 - y^2$  at  $(1, 2)$  in the direction of  $4\hat{i} + 3\hat{j}$  is

- (a)  $\frac{4}{5}$    (b)  $\frac{3}{5}$    (c)  $-\frac{4}{5}$    (d)  $-\frac{3}{5}$

Soln: $\rightarrow D.D. = \vec{\nabla} f \cdot \frac{\vec{a}}{|\vec{a}|} = (2x\hat{i} - 2y\hat{j}) \cdot \frac{(4\hat{i} + 3\hat{j})}{5} = \frac{8x - 6y}{5} \Big|_{(1,2)} = \frac{8 - 12}{5} = -\frac{4}{5}$

2. The directional derivative  $\phi = x^2 - y^2 + 2z^2$  at  $P(1, 2, 3)$  in direction  $\overrightarrow{PQ}$  where  $Q = (5, 0, 4)$  \_\_\_\_\_.

$$\text{Soln: } \overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = (5, 0, 4) - (1, 2, 3) = (4, -2, 1)$$

$$\begin{aligned} \therefore D.A. &= \nabla \phi \cdot \frac{\overrightarrow{PQ}}{|\overrightarrow{PQ}|} = (2x\hat{i} - 2y\hat{j} + 4z\hat{k}) \cdot \frac{(4\hat{i} - 2\hat{j} + \hat{k})}{\sqrt{16+4+1}} \\ &= \frac{8x + 4y + 4z}{\sqrt{21}} \Big|_{(1, 2, 3)} = \frac{28}{\sqrt{21}}. \end{aligned}$$

3. The directional derivative  $f = xy^2 + yz^2 + zx^2$  along to the tangent to the curve  $x=t$ ,  $y=t^2$  &  $z=t^3$  at  $(1, 1, 1)$  is \_\_\_\_\_.

Soln: Vector eqn of the curve is given by

$$\begin{aligned} \vec{x}(t) &= x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k} \\ &= t\hat{i} + t^2\hat{j} + t^3\hat{k} \end{aligned}$$

$$\frac{d\vec{x}}{dt} \Big|_{(1, 1, 1)} = \hat{i} + 2t\hat{j} + 3t^2\hat{k} \Big|_{t=1} = \hat{i} + 2\hat{j} + 3\hat{k} = \vec{a}$$


$$\begin{aligned} \therefore D.A. &= \nabla f \cdot \frac{\vec{a}}{|\vec{a}|} = [(y^2 + 2zx)\hat{i} + (2xy + z^2)\hat{j} + (2yz + x^2)\hat{k}] \cdot \frac{(\hat{i} + 2\hat{j} + 3\hat{k})}{\sqrt{1+4+9}} \\ &= \frac{18}{\sqrt{14}}. \end{aligned}$$

4. The directional derivative of  $\phi = \frac{y}{x^2+y^2}$  along the line which makes angle  $30^\circ$  with positive  $x$ -axis at  $(0, 1)$  is \_\_\_\_\_.

$$\text{Soln: } \vec{x} = x\hat{i} + y\hat{j}$$

$$\vec{x} = r\cos\theta\hat{i} + r\sin\theta\hat{j}$$

$$\Rightarrow \frac{\vec{x}}{r} = \cos\theta\hat{i} + \sin\theta\hat{j}$$



Note: The vector  $\vec{a}$  of a st. line which makes an angle  $\theta$  with +ve  $x$ -axis is  $\cos\theta\hat{i} + \sin\theta\hat{j}$ .

$$\therefore \frac{\vec{r}}{r} = \cos 30^\circ \hat{i} + \sin 30^\circ \hat{j} = \frac{\sqrt{3}}{2} \hat{i} + \frac{1}{2} \hat{j}$$

37

$$\therefore D.D. = \nabla \phi \cdot \frac{\vec{r}}{r} = -\frac{1}{2}$$

5. The greatest rate of increase of  $\phi = x^2yz$  at  $(2, -1, 2)$  is \_\_\_\_\_.

$$\text{Soln: } \nabla \phi = 2xyz \hat{i} + x^2z \hat{j} + x^2y \hat{k}$$

$$\therefore \nabla \phi \Big|_{(2, -1, 2)} = -8\hat{i} + 8\hat{j} - 4\hat{k}$$

Greatest rate of increase (or) max. value of D.D. =  $|\nabla \phi|$

$$= \sqrt{64+64+16}$$

$$= \sqrt{144} = 12$$

6. The angle b/w the surfaces  $x^2+y^2+z^2=9$  &  $x^2+y^2-z=3$  at  $(2, -1, 2)$  is \_\_\_\_\_.

$$\text{Soln: } \text{Let, } \phi_1 = x^2+y^2+z^2 \Rightarrow \nabla \phi_1 = 2x\hat{i} + 2y\hat{j} + 2z\hat{k}$$

$$\Rightarrow \nabla \phi_1 \Big|_{(2, -1, 2)} = 4\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\text{Let, } \phi_2 = x^2+y^2-z \Rightarrow \nabla \phi_2 = 2x\hat{i} + 2y\hat{j} - \hat{k}$$

$$\Rightarrow \nabla \phi_2 \Big|_{(2, -1, 2)} = 4\hat{i} - 2\hat{j} - \hat{k}$$

$$\cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|} = \frac{16+4-4}{\sqrt{16+4+16} \sqrt{16+4+1}} = \frac{16}{\sqrt{36} \sqrt{21}} = \frac{16}{6\sqrt{21}}$$

$$\Rightarrow \theta = \cos^{-1} \left( \frac{8}{3\sqrt{21}} \right).$$

7. If  $\vec{a}$  &  $\vec{b}$  are two arbitrary vectors with magnitudes  $a$  &  $b$  respectively

$$\text{then, } |\vec{a} \times \vec{b}|^2 = \underline{\hspace{2cm}}$$

$$\text{Soln: } \vec{a} \times \vec{b} = (\vec{a} \cdot \vec{b}) \sin \theta \hat{n}$$

$$|\vec{a} \times \vec{b}|^2 = a^2 b^2 \sin^2 \theta = a^2 b^2 (1 - \cos^2 \theta) = a^2 b^2 \left( 1 - \frac{(\vec{a} \cdot \vec{b})^2}{a^2 b^2} \right)$$

$$= a^2 b^2 - (\vec{a} \cdot \vec{b})^2$$

Divergence of a Vector function: → Let  $\vec{F}(x, y, z) = F_1 \hat{i} + F_2 \hat{j} + F_3 \hat{k}$  be

the differential vector point function then

$$\operatorname{div} \vec{F} = \vec{\nabla} \cdot \vec{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

Note: → If  $\vec{\nabla} \cdot \vec{F} = 0$  then  $\vec{F}$  is called solenoidal vector.

Curl of a Vector function: →

$$\operatorname{curl} \vec{F} = \vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix}$$

Note: → (i) If  $\vec{\nabla} \times \vec{F} = \vec{0}$  then  $\vec{F}$  is called irrotational vector.

(ii) If  $\vec{v} \rightarrow$  velocity vector

$\vec{\omega} \rightarrow$  angular velocity

then,  $\vec{v} = \vec{\omega} \times \vec{r}$

$$\operatorname{curl} \vec{v} = \vec{\nabla} \times (\vec{\omega} \times \vec{r}) = 2\vec{\omega}$$

$$\Rightarrow \boxed{\vec{\omega} = \frac{1}{2} \operatorname{curl} \vec{v}}$$

Scalar Potential function: → For every rotation<sup>al</sup> vector,  $\exists$  a function  $\phi$  scalar.

s.t.  $\vec{F} = \vec{\nabla} \phi$ , then  $\phi$  is said to be scalar potential fun.

Note: → (i)  $\operatorname{curl}(\operatorname{grad} \phi) = \vec{0}$

(ii)  $\operatorname{div}(\operatorname{curl} \vec{F}) = 0$

(iii)  $\operatorname{div}(\operatorname{grad} \phi) = \nabla(\nabla \phi) = \nabla^2 \phi$

where,  $\nabla^2 \rightarrow$  Laplacian operator

$$\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

Questions:-

1. The value of  $\nabla \cdot (\gamma^n \vec{r}) = \underline{\hspace{2cm}}$ , where  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

- (a)  $(n+3)\gamma^n$       (b)  $(n-2)\gamma^n$       (c)  $n\gamma^{n-3}$       (d)  $(n+2)\gamma^{n-1}$

and hence which of the following is solenoidal

(a)  $\gamma^3 \vec{r}$

(b)  $\gamma \vec{r}$

(c)  $\frac{\vec{r}}{\gamma^3}$

(d)  $\frac{\vec{r}}{\gamma^2}$

Sol<sup>n</sup>:  $\gamma^n \vec{r} = \underbrace{\gamma^n x \hat{i}}_{f_1} + \underbrace{\gamma^n y \hat{j}}_{f_2} + \underbrace{\gamma^n z \hat{k}}_{f_3}$

$$\nabla \cdot (\gamma^n \vec{r}) = \frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} + \frac{\partial f_3}{\partial z}$$

$$\begin{aligned}\frac{\partial f_1}{\partial x} &= \frac{\partial}{\partial x} (\gamma^n x) = \gamma^n + x n \gamma^{n-1} \frac{\partial \gamma}{\partial x} \\ &= \gamma^n + n x \gamma^{n-1} \frac{x}{\gamma} \\ &= \gamma^n + n x \gamma^{n-2} x^2\end{aligned}$$

Similarly,  $\frac{\partial f_2}{\partial y} = \gamma^n + n \gamma^{n-2} y^2, \quad \frac{\partial f_3}{\partial z} = \gamma^n + n \gamma^{n-2} z^2$

$$\begin{aligned}\therefore \nabla \cdot (\gamma^n \vec{r}) &= 3\gamma^n + n \gamma^{n-2} (x^2 + y^2 + z^2) = 3\gamma^n + n \gamma^{n-2} \cdot r^2 \\ &= 3\gamma^n + n \gamma^n = (n+3)\gamma^n\end{aligned}$$

Now, for solenoidal,  $\nabla \cdot (\gamma^n \vec{r}) = 0$

$$\Rightarrow (n+3)\gamma^n = 0 \Rightarrow n = -3$$

2.  $\nabla \cdot \left( \frac{\vec{r}}{\gamma^3} \right) = 0$ .

2. If  $\vec{F} = (3x^2 + 2y)\hat{i} - 4xz\hat{j} + 3xy^2\hat{k}$  represents a velocity vector then

corresponding angular velocity at  $(2, 2, -1)$  is \_\_\_\_\_.

Sol<sup>n</sup>:  $\text{curl } \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3x^2 + 2y & -4xz & 3xy^2 \end{vmatrix} = 32\hat{i} - 12\hat{j} + 2\hat{k}$

$\therefore \vec{\omega} = \frac{1}{2} \text{curl } \vec{F} = 16\hat{i} - 6\hat{j} + \hat{k}$

3. If  $\phi(x,y) = ax^2y - y^3$  &  $\nabla^2\phi = 0$  then,  $a = \underline{\hspace{2cm}}$ .

- (a) 2      (b) 3      (c) -2      (d) -3

$$\text{Soln: } \rightarrow \nabla^2\phi = \frac{\partial^2\phi}{\partial x^2} + \frac{\partial^2\phi}{\partial y^2} + \frac{\partial^2\phi}{\partial z^2} = 0$$

$$\Rightarrow \cancel{2ay} - 6y = 0 \Rightarrow a = 3.$$

4. If  $\vec{F} = (5x+7z^2)\hat{i} + (4x^2+\lambda y)\hat{j} + (7z-2xy)\hat{k}$  is solenoidal then  $\lambda = \underline{\hspace{2cm}}$

$$\text{Soln: } \rightarrow \vec{\nabla} \cdot \vec{F} = 0$$

$$\Rightarrow 5 + \lambda + 7 = 0 \Rightarrow \lambda = -12.$$

5. If  $\vec{F} = 5x^2z\hat{i} - 7xy^2\hat{j} + (12x+7z)\hat{k}$  then  $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{F})$  at  $(5, 3, -2)$  is 0.

6. If  $\vec{F} = (x+2y+az)\hat{i} + (bx-3y-z)\hat{j} + (4x+cy+2z)\hat{k}$  is irrotational vector

fun then  $a, b, c = \underline{\hspace{2cm}}$ .

$$\text{Soln: } \rightarrow \vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x+2y+az & bx-3y-z & 4x+cy+2z \end{vmatrix}$$

$$= \hat{i}(4-a) + \hat{j}(4-a) + \hat{k}(b-2)$$

$$= 0$$

$$\therefore a = 4, b = 2.$$

## 7. Vector Integration: $\rightarrow$

Line Integral: An integral evaluated over a curve is called line integral.

Let,  $\vec{F}(x, y, z) = f_1\hat{i} + f_2\hat{j} + f_3\hat{k}$  be a differentiable point ~~function~~

function defined at each point on the curve 'c' then its line integral is

$$\int_C \vec{F} \cdot d\vec{r} = \int_C (f_1 dx + f_2 dy + f_3 dz)$$

Note: If  $C$  is a closed curve, then line integral of  $\vec{F}$  over ' $C$ ' is called circulation of  $\vec{F}$  i.e.  $\oint_C \vec{F} \cdot d\vec{r}$ . 39

Work done by a force: → The total work done by a force  $\vec{F}$  in moving a particle along a curve ' $C$ ' is  $\int_C \vec{F} \cdot d\vec{r}$ .

Note: If  $\vec{F}$  is irrotational, then, the line integral of  $\vec{F}$  is independent of the path.

i.e. when  $\vec{F}$  is irrotational we have  $\vec{F} = \nabla \phi$ , where  $\phi$  is a scalar potential fn then,

$$\int_a^b \vec{F} \cdot d\vec{r} = \phi_b - \phi_a .$$

Questions: →

1. The value of  $\int_C \vec{F} \cdot d\vec{r}$ , where,  $\vec{F} = 3xy\hat{i} - y^2\hat{j}$  if ' $C$ ' is the curve  $y=2x^2$  joining pts  $(0,0)$  &  $(1,2)$  is \_\_\_\_\_.

$$\text{Soln: } \int_C \vec{F} \cdot d\vec{r} = \int_C (F_1 dx + F_2 dy + F_3 dz) = \int_C (3xy dx - y^2 dy)$$

$$= \int_0^1 [3x(2x^2) dx - 4x^4 \cdot 4x dx] \quad \left. \begin{array}{l} \Rightarrow y=2x^2 \\ \Rightarrow dy = 4x dx \end{array} \right\}$$


$$= \frac{6}{4} - \frac{16}{6} =$$

2. The value of  $\int_C \vec{F} \cdot d\vec{r}$ , where,  $\vec{F} = 3xy\hat{i} - y^2\hat{j}$  if ' $C$ ' is the curve bounded by  $y=x$  &  $y=x^2$  is \_\_\_\_\_.

$$\text{Soln: } \int_C \vec{F} \cdot d\vec{r} = \int_{C_1} + \int_{C_2}$$

along  $C_1$ ,  $y=x^2 \Rightarrow dy = 2x dx$

$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_1} (3xy dx - y^2 dy)$$



$$= \int_0^1 [3x(x^2) dx - x^4 \cdot 2x dx] = \frac{5}{12}$$

Now, along  $C_2$ ,  $y=x \Rightarrow dy=dx$

$$\therefore \int_{C_2} \vec{F} \cdot d\vec{x} = \int_1^0 [3x(x) dx - x^2 dx] = -\frac{2}{3}$$

$$\therefore \int \vec{F} \cdot d\vec{x} = \frac{5}{12} - \frac{2}{3} = -\frac{3}{12} = -\frac{1}{4}.$$

3. The value of  $\int_C [(3x+4y) dx + (2x-3y) dy]$  where, 'C' is the circle of radius 2, with centre at origin in x-y plane is \_\_\_\_\_.

- (a)  $4\pi$     (b)  $8\pi$     (c)  $-4\pi$     (d)  $-8\pi$

Soln: Here,  $C \rightarrow x^2+y^2=2^2$

whenever the curve is circle we will go for polar form.

$$\text{let, } x=2\cos\theta; y=2\sin\theta$$

$$\Rightarrow dx=-2\sin\theta d\theta, dy=2\cos\theta d\theta$$

$$\int_0^{2\pi} [(3x2\cos\theta + 4x2\sin\theta)(-2\sin\theta d\theta) + (2x2\cos\theta - 3x2\sin\theta)(2\cos\theta d\theta)]$$

$$= \int_0^{2\pi} [-12\sin\theta\cos\theta - 16\sin^2\theta d\theta + 8\cos^2\theta d\theta - 12\sin\theta\cos\theta d\theta]$$

$$= \int_0^{2\pi} [-24\sin\theta\cos\theta d\theta]$$

4. The value of  $\int_C \vec{F} \cdot d\vec{r}$ , where,  $\vec{F} = (2y+3)\hat{i} + xz\hat{j} + (yz-x)\hat{k}$  along the line joining

40

(i)  $(0,0,1)$  &  $(0,1,1)$

(ii)  $(0,1,1)$  &  $(2,1,1)$

is \_\_\_\_\_.

Soln: → (i) Along that st. line  $x=0, z=1 \Rightarrow dx=0, dz=0$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C F_2 dy = \int_0^1 xz dy = 0.$$

(ii)  $y=1, z=1 \Rightarrow dy=0, dz=0$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C F_1 dx = \int_0^2 (2y+3) dx = \int_0^2 5 dx = 5x \Big|_0^2 = 10.$$

5. The total work done by a force  $\vec{F} = (3x^2+6y)\hat{i} - 14yz\hat{j} + 20xz^2\hat{k}$  in moving a particle along a st. line joining the pts.  $(0,0,0)$  &  $(1,2,3)$

is \_\_\_\_\_.

Soln: →  $\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1} \Rightarrow \frac{x-0}{1-0} = \frac{y-0}{2-0} = \frac{z-0}{3-0} = t$

$$\Rightarrow x=t, y=2t, z=3t$$

$$dx=dt, dy=2dt, dz=3dt$$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C (F_1 dx + F_2 dy + F_3 dz)$$

$$= \int_0^1 [(3t^2+12t)dt + (-14 \times 2t \times 3t)(2dt) + (20x \times t \times 9t^2)(3dt)]$$

$$= \frac{540}{4}$$

6. The line integral  $\int \vec{F} \cdot d\vec{r}$  of fun  $\vec{F} = 2xyz\hat{i} + x^2z\hat{j} + x^2y\hat{k}$  from the origin  $(0,0,0)$  to the pt.  $(1,1,1)$  is \_\_\_\_.

- (a) 0      (b) 1      (c) -1

(d) cannot be determined without specifying the path.

$$\text{Soln: } \rightarrow \nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xyz & x^2z & x^2y \end{vmatrix} = \hat{i}(x^2 - x^2) - \hat{j}(2xy - 2xy) + \hat{k}(2xz - 2xz) = \vec{0}$$

$\Rightarrow F$  is irrotational.

$$\Rightarrow \vec{F} = \nabla \phi = \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k}$$

$$\Rightarrow \frac{\partial \phi}{\partial x} = 2xyz, \frac{\partial \phi}{\partial y} = x^2z, \frac{\partial \phi}{\partial z} = x^2y$$

The total differentiation of  $\phi$ ,

$$\begin{aligned} d\phi &= \frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz \\ &= 2xyz dx + x^2z dy + x^2y dz \\ &= d(x^2yz) \end{aligned}$$

$$\Rightarrow \phi = x^2yz$$

$$\therefore \int_a^b \vec{F} \cdot d\vec{r} = \phi_b - \phi_a = \phi_{(1,1,1)} - \phi_{(0,0,0)} = 1.$$

Green's Theorem in a Plane:  $\rightarrow$  Let  $M(x,y)$  &  $N(x,y)$  be continuous fun having continuous 1<sup>st</sup> order partial derivative defined in the closed region  $R$  bounded by the closed curve 'C' then,

$$\oint_C (M dx + N dy) = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy$$

Questions:-

1. Evaluate  $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$ , where,  $C$  is a curve bounded by  $x=0, y=0$  &  $x+y=1$ .

u1


$$\text{Soln: } \rightarrow M = 3x^2 - 8y^2$$

$$\Rightarrow \frac{\partial M}{\partial y} = -16y$$

$$N = 4y - 6xy$$

$$\Rightarrow \frac{\partial N}{\partial x} = -6y$$

$$\therefore \frac{\partial M}{\partial x} - \frac{\partial N}{\partial y} = 10y$$



$$\therefore \oint_C M dx + N dy = \iint_R \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dx dy$$


$$= \iint_0^{1-x} 10y \, dx \, dy = 5 \int_0^1 (1-x)^2 \, dx = \frac{5}{3}.$$

2. Evaluate  $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$ , where,  $C$  is a curve bounded by  $y=\sqrt{x}$  &  $y=x^2$ .

$$\text{Soln: } \rightarrow M = 3x^2 - 8y^2$$

$$\oint_C M dx + N dy = \iint_0^1 \frac{1}{x} 10y \, dx \, dy$$


$$= 5 \int_0^1 (x - x^4) \, dx = \frac{3}{2}.$$



3. Evaluate  $\oint_C xy \, dy - y^2 \, dx$ , where,  $C$  is a square cut from the first quadrant from by the lines  $x=1$  &  $y=1$ .

$$\text{Soln: } \rightarrow M = -y^2 \Rightarrow \frac{\partial M}{\partial y} = -2y$$

$$N = xy \Rightarrow \frac{\partial N}{\partial x} = y$$



$$\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} = 3y$$

$$\therefore \oint_C M dx + N dy = \iint_0^1 3y \, dx \, dy = \frac{3}{2}.$$

Surface Integral: → Let  $\vec{F}(x, y, z) = f_1 \hat{i} + f_2 \hat{j} + f_3 \hat{k}$  be differentiable vector point fun defined over the surface  $S$  then, its surface integro

$$\text{is } \int_S \vec{F} \cdot d\vec{s} = \int_S \vec{F} \cdot \vec{n} \, ds$$

where,  $\vec{n}$  → unit outward drawn normal to the surface.

In cartesian form,

$$\int_S \vec{F} \cdot d\vec{s} = \int_S \vec{F} \cdot \vec{n} \, ds = \int (f_1 dy dz + f_2 dx dz + f_3 dx dy)$$

Methods of Evaluation:

(i) If  $R_1$  is the projection of 'S' on to x-y plane then,

$$\int_S \vec{F} \cdot \vec{n} \, ds = \iint_{R_1} \vec{F} \cdot \vec{n} \frac{dx dy}{|\vec{n} \cdot \vec{k}|}$$

(ii) If  $R_2 \rightarrow y-z$  plane then,

$$\int_S \vec{F} \cdot \vec{n} \, ds = \iint_{R_2} \vec{F} \cdot \vec{n} \frac{dy dz}{|\vec{n} \cdot \vec{i}|}$$

(iii) If  $R_3 \rightarrow x-z$  plane then,

$$\int_S \vec{F} \cdot \vec{n} \, ds = \iint_{R_3} \vec{F} \cdot \vec{n} \frac{dx dz}{|\vec{n} \cdot \vec{j}|}$$

Questions:-

42

1. The value of  $\int_S \vec{F} \cdot \vec{n} ds$ , where,  $\vec{F} = z\hat{i} + x\hat{j} - 3y^2z\hat{k}$  and S is the surface of the cylinder  $x^2 + y^2 = 16$  included in the 1<sup>st</sup> octant between  $z=0$  &  $z=5$ .

Soln:- Let  $\phi = x^2 + y^2$

$$\vec{\nabla}\phi = 2x\hat{i} + 2y\hat{j}$$

$$\vec{n} = \frac{\vec{\nabla}\phi}{|\vec{\nabla}\phi|} = \frac{2x\hat{i} + 2y\hat{j}}{\sqrt{4x^2 + 4y^2}} = \frac{x\hat{i} + y\hat{j}}{\sqrt{x^2 + y^2}}$$

$$\vec{n} = \frac{x\hat{i} + y\hat{j}}{4}$$

$$\vec{F} \cdot \vec{n} = \frac{xz}{4} + \frac{xy}{4} = \frac{x}{4}(y+z)$$

Let, R  $\rightarrow$  y-z plane.

$$\begin{aligned} \int_S \vec{F} \cdot \vec{n} ds &= \iint_R \vec{F} \cdot \vec{n} \cdot \frac{dydz}{|\vec{n}| \cdot \hat{i}} = \iint_R \frac{x}{4}(y+z) \cdot \frac{dydz}{(x/4)} \\ &= \int_{z=0}^5 \int_{y=0}^4 (y+z) dy dz \\ &= \int_0^5 \left(\frac{y^2}{2} + yz\right)_0^4 dz = 90. \end{aligned}$$

2. The value of  $\int_S \vec{F} \cdot \vec{n} ds$ , where,  $\vec{F} = 4xz\hat{i} - y^2\hat{j} + yz\hat{k}$  & S is surface of the cube bounded by  $x=0, x=1, y=0, y=1, z=0$  &  $z=1$ , is \_\_\_\_\_

Soln:- Method 1:-


$$\int_S \vec{F} \cdot \vec{n} ds = \int_{S_1} + \int_{S_2} + \dots + \int_{S_6}$$

Over  $S_1$ :- In x-y plane (OAPR);

$$z=0, \vec{n} = -\hat{k}$$

$$\vec{F} \cdot \vec{n} = -yz = 0$$

$$\therefore \int_{S_1} \vec{F} \cdot \vec{n} ds = 0$$



Over  $S_2$ : - parallel to  $xy$  plane (SQRc)

$$z=1, \vec{n} = \hat{k}, \vec{F} \cdot \vec{n} = yz = y$$

$$\therefore \int_{S_2} \vec{F} \cdot \vec{n} ds = \iint_R \vec{F} \cdot \vec{n} \frac{dx dy}{|\vec{n} \cdot \hat{k}|} = \iint_0^1 y dx dy = \frac{1}{2}$$

Over  $S_3$ : - in  $y-z$  plane (OBRC)

$$x=0, \vec{n} = -\hat{i}, \vec{F} \cdot \vec{n} = -4xz = 0$$

$$\therefore \int_{S_3} \vec{F} \cdot \vec{n} ds = 0$$

over  $S_4$ : - parallel to  $y-z$  plane (AP)

$$x=1, \vec{n} = \hat{i}, \vec{F} \cdot \vec{n} = 4xz = 4z$$

$$\therefore \int_{S_4} \vec{F} \cdot \vec{n} ds = \iint_R \vec{F} \cdot \vec{n} \frac{dy dz}{|\vec{n} \cdot \hat{i}|} = \iint_0^1 4z dy dz = 2$$

over  $S_5$ : - in  $x-z$  plane (OCSA)

$$y=0, \vec{n} = -\hat{j}, \vec{F} \cdot \vec{n} = y^2 = 0$$

$$\therefore \int_{S_5} \vec{F} \cdot \vec{n} ds = 0$$

over  $S_6$ : - parallel to  $x-z$  plane (BRQP)

$$y=1, \vec{n} = \hat{j}, \vec{F} \cdot \vec{n} = -y^2 = -1$$

$$\therefore \int_{S_6} \vec{F} \cdot \vec{n} ds = \iint_R \vec{F} \cdot \vec{n} \frac{dx dz}{|\vec{n} \cdot \hat{j}|} = \iint_0^1 (-1) dx dz = -1$$

~~$$\therefore \int_S \vec{F} \cdot \vec{n} ds = 0 + \frac{1}{2} + 0 + 2 + 0 - 1 = \frac{3}{2}$$~~

Method 2:

$$\int_S \vec{F} \cdot \vec{n} ds = \iint_V \nabla \cdot \vec{F} dV \quad (\text{Divergence Theorem})$$

$$\nabla \cdot \vec{F} = 4z - 2y + y = 4z - y$$

$$\therefore \int_S \vec{F} \cdot \vec{n} ds = \iiint_0^1 (4z - y) dz dy dx = \frac{3}{2}$$

43

Volume Integral: Let  $\vec{F}(x, y, z) = f_1\hat{i} + f_2\hat{j} + f_3\hat{k}$  be the differentiable vector point fun defined in volume  $V$ , then, its volume integral is  $\int_V \vec{F} \cdot dV$

if  
Similarly,  $\phi(x, y, z)$  is scalar point fun, then,  $\int \phi dV$

Questions:

- Evaluate  $\int_V \vec{F} \cdot dV$  where,  $\vec{F} = (2x^2 - 3z)\hat{i} - 2xy\hat{j} - 4x\hat{k}$  and  $V$  is the region bounded by the planes  $x=0, y=0, z=0$  &  $2x+2y+z=4$ .

Soln:  $\vec{\nabla} \cdot \vec{F} = 4x - 2x = 2x$

$$\int 2x dV = \int \int \int_{x=0, y=0, z=0}^{2x, 4-2x-2z} 2x dz dy dx = \frac{8}{3}$$

- The volume of an object expressed in spherical coordinate,  $V = \int \int \int_{0, 0, 0}^{2\pi, \pi/2, 1} r^2 \sin\theta dr d\phi d\theta$ , then  $V$  is \_\_\_\_\_.

- (a)  $\frac{\pi}{3}$  (b)  $\frac{\pi}{6}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{\pi}{4}$

Soln:  $V = \int \int \int_{0, 0, 0}^{2\pi, \pi/2, 1} \sin\theta \frac{r^2}{2} dr d\phi d\theta = \frac{1}{2} \int_{0}^{2\pi} (-\cos\theta) \Big|_{0}^{\pi/2} d\theta = \frac{\pi}{3}$

Gauss-Divergence Theorem: Let  $S$  be a closed surface enclosing a volume  $V$  &  $\vec{F}(x, y, z) = f_1\hat{i} + f_2\hat{j} + f_3\hat{k}$  be the differentiable vector point fun defined over  $S$ , then,

$$\int_S \vec{F} \cdot d\vec{s} = \int_V \operatorname{div} \vec{F} dV$$

Questions:

- Evaluate  $\int_S \vec{F} \cdot \vec{n} ds$ , where,  $\vec{F} = x\hat{i} + y\hat{j} + z\hat{k}$  &  $S$  is a closed surface

enclosing a volume  $V$  is \_\_\_\_\_.

- (a)  $V$  (b)  $2V$  (c)  $3V$  (d)  $4V$

2. The value of  $\int_S (x \, dy \, dz + y \, dx \, dz + z \, dx \, dy)$ , where  $S$  is a surface,

(i) cylinder  $x^2 + z^2 = 16$  &  $y=0$  to  $y=3$ ,

(ii) is a sphere  $x^2 + y^2 + z^2 = 9$ ,

is \_\_\_\_\_.

$$\text{Soln: } \vec{F} = x\hat{i} + y\hat{j} + z\hat{k} \quad (\because \int_S \vec{x} \cdot \vec{n} \, ds = \int_S \vec{F} \cdot d\vec{s} = \int_S (F_1\hat{i} + F_2\hat{j} + F_3\hat{k}) \, d\vec{s})$$

$$\int_S \vec{x} \cdot \vec{n} \, ds = 3V$$

$$(i) \int_S \vec{x} \cdot \vec{n} \, ds = 3\pi r^2 h = 3\pi \times 4^2 \times 3 =$$

$$(ii) \int_S \vec{x} \cdot \vec{n} \, ds = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi \times 3^3 =$$

3. Evaluate  $\int_S (x^2 + 2y^2 + 3z^2) \, ds$ , where,  $S$  is the surface  $x^2 + y^2 + z^2 = 1$ .

$$\text{Soln: } \vec{F} \cdot \vec{n} = x^2 + 2y^2 + 3z^2$$

$$\phi = x^2 + y^2 + z^2$$

$$\vec{n} = \frac{\nabla \phi}{|\nabla \phi|} = x\hat{i} + y\hat{j} + z\hat{k}$$

Let,  $F_1\hat{i} + F_2\hat{j} + F_3\hat{k} = \vec{F}$ , then,

$$\vec{F} \cdot \vec{n} = F_1 x + F_2 y + F_3 z = x^2 + 2y^2 + 3z^2$$

$$\Rightarrow F_1 = x, F_2 = 2y, F_3 = 3z$$

$$\therefore \vec{F} = x\hat{i} + 2y\hat{j} + 3z\hat{k}$$

$$\therefore \int_S \vec{F} \, ds = \int_V \vec{\nabla} \cdot \vec{F} \, dv = \int_V (1+2+3) \, dv = 6V = \frac{6 \times 4}{3} \pi \times 1 = 8\pi.$$

4. The value of  $\int_S \vec{F} \cdot \vec{n} \, ds$ , where  $\vec{F} = 4x^2\hat{i} - 3y\hat{j} + 8xz\hat{k}$  &  $S$  is a surface  $0 \leq x \leq 1, 0 \leq y \leq 2, \& 0 \leq z \leq 3$ , is \_\_\_\_\_.

$$\text{Soln: } \vec{\nabla} \cdot \vec{F} = 8x - 3 + 8x = 16x - 3$$

$$\therefore \int_S \vec{F} \cdot \vec{n} \, ds = \int_V \vec{\nabla} \cdot \vec{F} \, dv = \int_{x=0}^1 \int_{y=0}^2 \int_{z=0}^3 (16x - 3) \, dx \, dy \, dz = 30.$$

5. Evaluate  $\int_S \vec{F} \cdot \vec{n} ds$ , where,  $\vec{F} = 4x\hat{i} - 2y^2\hat{j} + z^2\hat{k}$ , taken over the region bounded by  $x^2 + y^2 = 4$  &  $z=0$  to  $z=3$ .

$$\text{Soln: } \rightarrow \vec{\nabla} \cdot \vec{F} = 4 - 4y + 2z$$

$$\begin{aligned}\therefore \int_S \vec{F} \cdot \vec{n} ds &= \int_V \vec{\nabla} \cdot \vec{F} dV = \int_{x=-2}^2 \int_{y=-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{z=0}^{3} (4 - 4y + 2z) dx dy dz \\ &= 2 \int_0^3 \int_{-2}^2 (4 + 2z) \sqrt{4-x^2} dx dz \\ &= 4 \int_0^3 \int_0^2 (4 + 2z) \sqrt{4-x^2} dx dz \\ &= 4 \int_0^3 (4 + 2z) \left[ \frac{x\sqrt{4-x^2}}{2} + \frac{4}{2} \sin^{-1}\left(\frac{x}{2}\right) \right]_0^2 dz \\ &= 4\pi \int_0^3 (4 + 2z) dz = 84\pi\end{aligned}$$

Note: → we can also solve the above problem in polar form.

6. Evaluate  $\int_S \vec{\nabla} \times \vec{F} \cdot \vec{n} ds$ , where,  $\vec{F} = 4x^2z\hat{i} - (yz - 7)\hat{j} + xy^2z\hat{k}$ , and  $S$  is the surface bounded by  $y^2 + z^2 = 25$  &  $x=0$  to  $x=2$ .

$$\text{Soln: } \rightarrow \int_S \vec{\nabla} \times \vec{F} \cdot \vec{n} ds = \int_V \operatorname{div}(\vec{\nabla} \times \vec{F}) dV = 0.$$

Stoke's Theorem: → Let  $S$  be an open surface bounded by a closed curve  $C$  &  $\vec{F}(x, y, z) = f_1\hat{i} + f_2\hat{j} + f_3\hat{k}$  be a differentiable vector function defined over ' $S$ ', then  $\oint_C \vec{F} \cdot d\vec{s} = \int_S \vec{\nabla} \times \vec{F} \cdot d\vec{s} = \int_S \vec{\nabla} \times \vec{F} \cdot \vec{n} ds$

$$\text{i.e. } \oint_C (f_1 dx + f_2 dy + f_3 dz) = \int_S (\vec{\nabla} \times \vec{F}) \cdot \vec{n} ds$$

Questions: → 1. The value of  $\int_C \vec{F} \cdot d\vec{x}$ , where  $\vec{F} = yz\hat{i} + xz\hat{j} + xy\hat{k}$ , if 'C' is a curve  $x^2 + y^2 = 4$  in XY plane is \_\_\_\_\_.

- (a) 0    (b)  $\frac{1}{2}$     (c) 2    (d) 3

Sol<sup>n</sup>: →  $\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ yz & xz & xy \end{vmatrix} = \hat{i}(x-x) - \hat{j}(y-y) + \hat{k}(z-z) = \vec{0}$

$$\therefore \int_C \vec{F} \cdot d\vec{x} = \int_S (\vec{\nabla} \times \vec{F}) \cdot \vec{n} ds = \int_S \vec{0} \cdot \vec{n} ds = 0.$$

2. The value of  $\int_C \vec{F} \cdot d\vec{x}$ , where  $\vec{F} = -y^3\hat{i} + x^3\hat{j}$  & 'C' is the circular disc  $x^2 + y^2 \leq 1$ ,  $z=0$  is \_\_\_\_\_.

Sol<sup>n</sup>: →  $\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -y^3 & x^3 & 0 \end{vmatrix} = \hat{i}(0-0) - \hat{j}(0-0) + \hat{k}(3x^2 + 3y^2) = 3(x^2 + y^2)\hat{k}$

$$\therefore \vec{\nabla} \times \vec{F} \cdot \vec{n} = 3(x^2 + y^2) \quad \left\{ \because \vec{n} = \hat{k} \right\}$$

$$\therefore \int_C \vec{F} \cdot d\vec{x} = \int_S (\vec{\nabla} \times \vec{F}) \cdot \vec{n} ds$$

$$= \int_S 3(x^2 + y^2) ds$$

Let,  $R \rightarrow$  XY plane

$$\therefore \int_C \vec{F} \cdot d\vec{x} = \iint_R 3(x^2 + y^2) \frac{dx dy}{|\vec{n} \cdot \hat{k}|} = \iint_R 3(x^2 + y^2) dx dy$$

Let  $x = r \cos \theta$ ,  $y = r \sin \theta$

$$x^2 + y^2 = r^2, \quad |J| = r$$

$$\therefore \int_C \vec{F} \cdot d\vec{x} = \int_0^{2\pi} \int_0^1 3r^2 r dr d\theta = \frac{3\pi}{2}.$$

3. Evaluate  $\oint_C (y dx + z dy + x dz)$ , where  $C$  is the curve of intersection

$$\text{of } x^2 + y^2 + z^2 = a^2 \text{ & } x+z=a.$$

Soln:  $\vec{F} = y\hat{i} + z\hat{j} + x\hat{k}$

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y & z & x \end{vmatrix} = \hat{i}(0-1) - \hat{j}(1-0) + \hat{k}(0-1) \\ = -\hat{i} - \hat{j} - \hat{k}$$

The intersection of sphere  $x^2 + y^2 + z^2 = a^2$  with the plane  $x+z=a$

is a circle in the plane  $x+z=a$ , with AB as diameter, where

$$A(a, 0, 0) \text{ & } B(0, 0, a)$$

$$AB = \sqrt{a^2 + a^2} = a\sqrt{2}$$

$$\Rightarrow \text{radius} = \frac{a}{\sqrt{2}}$$

$$\text{Let, } \phi = x+z$$


$$\vec{\nabla} \phi = \hat{i} + \hat{k}$$

$$\vec{n} = \frac{\vec{\nabla} \phi}{|\vec{\nabla} \phi|} = \frac{\hat{i} + \hat{k}}{\sqrt{2}}$$

$$(\vec{\nabla} \times \vec{F}) \cdot \vec{n} = (-\hat{i} - \hat{j} - \hat{k}) \cdot \frac{(\hat{i} + \hat{k})}{\sqrt{2}} = -\sqrt{2}$$

$$\oint_C (y dx + z dy + x dz) = \int_S (\vec{\nabla} \times \vec{F}) \cdot \vec{n} ds = \int_S -\sqrt{2} ds = -\sqrt{2} S$$

$$= -\sqrt{2} \pi \times \left(\frac{a}{\sqrt{2}}\right)^2 = -\frac{\pi a^2}{\sqrt{2}}$$



B. Fourier Series : $\rightarrow$  Let  $f(x)$  be a periodic fun defined in  $(c; c+2l)$  with period  $2l$ , then, the fourier series of  $f(x)$  is given by,

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos \frac{n\pi x}{l} + b_n \sin \frac{n\pi x}{l}]$$

where,  $a_0$ ,  $a_n$  &  $b_n$  are fourier coeff. given by,

$$a_0 = \frac{1}{l} \int_c^{c+2l} f(x) dx$$

$$a_n = \frac{1}{l} \int_c^{c+2l} f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_c^{c+2l} f(x) \sin \frac{n\pi x}{l} dx$$

Note: $\rightarrow$   $[-l, l]$ ,  $[0, 2l]$ ,  $[-\pi, \pi]$  (or)  $[0, 2\pi]$

Airchilet's Conditions: $\rightarrow$  A. fun is said to satisfy dirichilet's cond if

- (i)  $f(x)$  and its integrals are finite & single valued.
- (ii)  $f(x)$  has finite no. of finite discontinuities.
- (iii)  $f(x)$  has finite no. of maxima & minima.

Note: $\rightarrow$  If  $f(x)$  satisfies Dirichlet's cond then the fourier series is convergent.

Convergence: $\rightarrow$  (i) If  $f(x)$  is continuous at  $x=c \in (a, b)$  then fourier series of  $f(x)$  at  $x=c$  converges to  $f(c)$ .

(ii) If  $f(x)$  is discontinuous at  $x=c \in (a, b)$  then fourier series of  $f(x)$

$$\text{at } x=c \text{ converges to } \frac{1}{2} [\lim_{x \rightarrow c^-} f(x) + \lim_{x \rightarrow c^+} f(x)]$$

(iii) fourier series of  $f(x)$  at the end ptz i.e.  $x=a$  (or)  $b$  converges to  $\frac{1}{2} [\lim_{x \rightarrow a^+} f(x) + \lim_{x \rightarrow b^-} f(x)]$

## Fourier Series of Even & Odd function in $[-l, l]$ or $[-\pi, \pi]$ :- 46

### (i) Fourier Series of an even function :-

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l}$$

$$\text{where, } a_0 = \frac{2}{l} \int_0^l f(x) dx$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$$

### (ii) Fourier Series of an Odd function :-

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\text{where, } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

## Half-Range Series :-

### (i) Half-Range cosine series in $[0, l]$ :-

$$F(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l}$$

$$\text{where, } a_0 = \frac{2}{l} \int_0^l f(x) dx$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$$

### (ii) Half-Range sine series in $[0, l]$ :-

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\text{where, } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

Questions: →

1. The coeff. of  $\sin x$  in the fourier series expansion of  $f(x) = x^2$ , in  $(-\pi, \pi)$  is —

(a)  $\sum \frac{(-1)^n}{n^2}$  (b)  $\sum \frac{1}{n^2}$  (c)  $\frac{\pi^2}{6}$  (d) 0

Sol<sup>n</sup>: → Even fun<sup>n</sup> hence, coeff. of  $\sin x = 0$ .

2. If  $f(x) = \begin{cases} 0 & ; -2 < x < 0 \\ 1 & ; 0 < x < 2 \end{cases}$

then the term independent of  $x$  in the fourier series of  $f(x)$  is —

(a) 0 (b) 1 (c)  $\frac{1}{2}$  (d) 2

Sol<sup>n</sup>: → Here,  $(-2, 2) \Rightarrow l = 2$

$$a_0 = \frac{1}{l} \int_{-l}^{l} f(x) dx = \frac{1}{2} \int_{-2}^{2} 1 \cdot dx = 1$$

$$\therefore \text{Independent term} = \frac{a_0}{2} = \frac{1}{2}$$

3. The fun  $f(x) = \begin{cases} -x+1 & ; -\pi \leq x \leq 0 \\ x+1 & ; 0 \leq x \leq \pi \end{cases}$

then  $f(x)$  has following terms in its expansion

(a) cosine (b) sine (c) both (d) cannot be determined

Sol<sup>n</sup>: →  $f(-x) = \begin{cases} x+1 & ; -\pi \leq -x \leq 0 \\ -x+1 & ; 0 \leq -x \leq \pi \end{cases}$

$$= \begin{cases} x+1 & ; \pi \geq x \geq 0 \\ -x+1 & ; 0 \geq x \geq -\pi \end{cases}$$

$$\therefore f(-x) = -f(x)$$

⇒ Even fun.

Q. 4: If  $f(x) = \begin{cases} 0 & -2 < x < 0 \\ 1 & 0 < x < 2 \end{cases}$

42

then the coeff. of  $\cos \frac{n\pi x}{2}$  is \_\_\_\_.

- (a)  $0$       (b)  $\frac{1}{n}$       (c)  $\frac{1}{n^2}$       (d)  $-\frac{1}{n}$

Sol<sup>n</sup>:  $\rightarrow (-2, 2) \Rightarrow l=2$

$$a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$$

$$= \frac{1}{2} \int_0^2 \cos \frac{n\pi x}{2} dx$$

$$= \frac{1}{2} \left[ \frac{\sin \frac{n\pi x}{2}}{\frac{n\pi}{2}} \right]_0^2$$

$$\approx \frac{1}{2} \times \frac{2}{n\pi} [0 - 0] = 0$$

5. If  $f(x) = x^2$  in  $[-\pi, \pi]$  has its fourier expansion as  $f(x) = \frac{\pi^2}{3} +$

$$4 \left[ \frac{\cos x}{1^2} - \frac{\cos 2x}{2^2} + \frac{\cos 3x}{3^2} - \dots \right] \text{ then the value of}$$

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots \text{ is } \dots$$

- (a)  $\frac{\pi^2}{6}$       (b)  $\frac{\pi^2}{12}$       (c)  $\pi^2$       (d) None


Sol<sup>n</sup>: At  $x = \pi$ , we have

$$\frac{\pi^2}{3} - 4 \left[ \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots \right] = \frac{1}{2} \left[ \lim_{x \rightarrow \pi^+} f(x) + \lim_{x \rightarrow \pi^-} f(x) \right]$$

$$= \frac{1}{2} [\pi^2 + \pi^2] = \pi^2$$

$$\Rightarrow \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{1}{4} \left[ \pi^2 - \frac{\pi^2}{3} \right] = \frac{\pi^2}{6}$$

6. A fun with period  $2\pi$  is shown below



then fourier series is \_\_\_\_\_

$$(a) f(x) = \sum_{n=1}^{\infty} \frac{2}{n\pi} \sin \frac{n\pi}{2} \cos nx$$

$$(b) f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n\pi} \sin \frac{n\pi}{2} \sin nx$$

$$\checkmark (c) f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2}{n\pi} \sin \frac{n\pi}{2} \cos nx$$

$$(d) f(x) = \sum_{n=1}^{\infty} \frac{2}{n\pi} \sin \frac{n\pi}{2} \sin nx$$

Soln: →

$$f(x) = \begin{cases} 0 &; -\pi < x < -\pi/2 \\ 1 &; -\pi/2 < x < \pi/2 \\ 0 &; \pi/2 < x < \pi \end{cases}$$

$f(-x) = f(x) \Rightarrow$  Even fun  $\Rightarrow$  cosine series

$$a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi/2} 1 dx = 1$$

$$\therefore \frac{a_0}{2} = 1$$

7. In  $[0, \pi]$  the constant term in the cosine series of  $f(x) = x^2 + 2x$  is

- (a)  $\pi(\frac{\pi}{3} - 1)$    (b)  $\pi(\frac{2\pi}{3} + 1)$    (c)  $\pi(\frac{\pi}{2} + 1)$    (d) None

Soln: →  $a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} [x^2 + 2x] dx = \frac{2}{\pi} \left[ \frac{x^3}{3} + x^2 \right]$

$$= \frac{2\pi^2}{3} + \pi^2$$

$$= \pi \left( \frac{2\pi^2}{3} + 1 \right)$$

$$\therefore \frac{a_0}{2} = \frac{1}{\pi} \left( \frac{\pi^3}{3} + \pi^2 \right) = \pi \left( \frac{\pi}{3} + 1 \right).$$

8. If  $f(x) = x$  is expressed on a half range cosine series in  $[0, 2]$   
then the coeff. of  $\cos \pi x$  is \_\_\_\_.

C18

- (a)  $\frac{4}{\pi^2}$  (b)  $\frac{2}{\pi^2}$  (c) 0 (d) None.

9. In the interval  $[0, \pi]$  if a const.  $c$  is expressed as a half range sine series then coeff. of  $\sin 5x$  is \_\_\_\_.

- (a)  $\frac{2c}{5\pi}$  (b) 0 (c)  $\frac{4c}{5\pi}$  (d)  $\frac{c}{5\pi}$

Sol<sup>n</sup>:  $\rightarrow f(x) = \sum_{n=1}^{\infty} b_n \sin nx, \quad f(x) = c$

$$b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cdot \sin nx dx$$

$$= \frac{2}{\pi} \int_0^{\pi} c \cdot \sin 5x dx$$

$$= \frac{2c}{\pi} \left[ -\frac{\cos 5x}{5} \right]_0^{\pi}$$

$$= \frac{4c}{5\pi}$$

